

Biotechnology NEWS



The Australian Stem Cell Summit
29 & 30 March 2007
Hilton on the Park, Melbourne
Workshops: Wednesday 28th
www.informa.com.au/s

[Editorial](#) | [Advertising](#) | [Privacy](#) | [About Us](#) | [Subscribe to Australian Life Scientist Magazine](#)
[Home](#) | [Biomedical](#) | [AgBiotech](#) | [Bioinformatics](#) | [Australian Life Scientist](#) | [Search](#)

Aligning biotechnology with business outcomes

Friday, 2nd February

SECTIONS

[Home](#)
[AgBiotech](#)
[Biomedical](#)
[Bioinformatics](#)

[Australian Life Scientist](#)

LATEST

[News](#)
[Features](#)
[Interviews](#)
[Market Reports](#)

TOPIC**CENTRES**

[Cloning](#)
[Drugs](#)
[Genetics](#)
[Genome](#)
[IPO](#)
[Proteome](#)
[Regulation](#)

DIGITAL**EDITIONS**[View latest](#)**ABOUT**

[Editorial](#)
[Advertising](#)
[Privacy](#)
[About Us](#)
[IDG Online](#)
[Contacts](#)

From laboratory to bedside

FIONA WYLIE, AUSTRALIAN LIFE SCIENTIST

15/11/2006 14:56:28

"It is not that stem cell transplantation doesn't work, it is just that we need more work to figure it out." With this kind of simple optimism, and a little green jasmine tea, Professor Brent Reynolds chatted with Fiona Wylie about life, coincidence and the use of neural stem cells to treat spinal cord injury.

Brian Reynolds is one of a distinguished list of speakers making up a two-part session, "In the search for a cure for spinal cord injury - from laboratory to bedside", at the Australian Health & Medical Research Congress (AH&MRC) at the Melbourne Convention Centre from November 26 to December 1.

Reynolds moved from Canada to the Queensland Brain Institute (QBI) at the University of Queensland in 2004. His path to this point has been somewhat unorthodox to say the least, particularly for someone who published a Science paper and devised an important new tool for the entire field during his PhD.

Immediately after finishing his doctorate in 1994, Reynolds founded a company called NeuroSpheres, based on this new technology.


"I was the director of research and we worked with large pharma and several biotechnology companies to further develop and protect the technology," Reynolds says. "Today NeuroSphere transplantation technology is licensed to Stem Cell Inc, based in California, who are about to start clinical trials based on technology we developed and patented, which is kind of exciting."

Impressively, the technology is also the basis of Phase II trials by another company for treating stroke, and at least half a dozen clinical trials starting in 2007-2008.

The unorthodox route to the QBI began in 1997, when Reynolds opted out of science to study Chinese medicine. He and his family spent the next few years between Thailand, running a yoga centre, and Salt Spring Island off the west coast of Canada, where Reynolds had a Chinese medicine clinic.

The lure back to science came in 2002 when an old university friend in Vancouver, who was head of business development with a company called StemCell Technologies, contacted him because the company wanted to get into the neural stem cell field.

"Things weren't working, he heard that I had moved to the west coast so he asked if I

**Jobs**

Looking for Staff
Advertise here. (Ric Hudson

Events

Holding an event
Advertise here. (Ric Hudson

LoginUsername: Password:

- If you have forgotten your password click here
- If you would like to subscribe click here

Advanced Search

Articles from:

2 2

Article to:

2 2

Keywords:

would come and have a look at this stuff, and I started going to help him one day a week," Reynolds says. "It was also near a really good yoga teacher."

Search

To help out, Reynolds got on the phone to former contacts looking for technology to license. One of these was Rietze, who used to work with Reynolds at NeuroSpheres. Rietze had just moved from Melbourne up to Queensland with Professor Perry Bartlett to set up the QBI.

It seems the next step was meant to be - Reynolds and his family had just been convinced by a friend from Thailand that Australia, and particularly Brisbane, was a great place to live.

"So Rod arrives and is telling me all about this new institute and says I should come and work with them in Brisbane. It was perfect."

Neural stem cells

Since his arrival in the Sunshine State, Reynolds and his team at the QBI have been developing methodology (which has been patented) to identify and expand distinct cell populations within a heterogeneous milieu of neural stem cell culture to benefit transplantation therapies, in particular spinal cord injury.

"The existence of stem cells in the adult mammalian central nervous system (CNS) and our ability to isolate and expand them ex vivo provides a number of therapeutic opportunities when it comes to treating spinal cord injury," Reynolds says.

Cell transplants into nervous tissue have been going on in animal models now for two to three decades. Primarily, the work has been done in rodent models of Parkinson's disease, with over 1000 reported studies of transplants into the brain. Human studies have also been carried out, with 300 to 400 people receiving human foetal tissue.

Basically, there were lots of promising results, and some not so promising results. Reynolds uses this history to highlight the two main problems with neural cell transplantation, which he will discuss along with ways to solve these problems at the AH&MRC.

Firstly, there is never going to be enough primary foetal tissue available for transplants, especially given the ethical and moral issues to be considered, he says. The primary requirement in this field is therefore a renewable source of stem cells.

One possibility is embryonic stem (ES) cells, differentiated down the neural lineage and grown up in culture for transplants. The problem with ES cells is that they are undifferentiated cells to start with and there is a chance, albeit only slight, that one of those cells doesn't terminally differentiate and grows to form a tumour after transplantation.

"All you are going to need is one tumour in one patient, and it will kill the whole field," he says. "That is what happened in the late '90s with gene therapy."

This issue is highlighted by a paper just published in Nature Medicine showing that human ES cells differentiated into dopamine neurons and transplanted into a rodent model of Parkinson's cured the symptoms of the disease. In the animals, however, developed tumours. The solution is better ways to sort cell populations early on in the process. Reynolds' group is also working on assays to do this.

The other possible renewable source are neural stem cells grown as neurospheres, which are clusters of cells in tissue culture from primary neural stem cells isolated from either adult or foetal tissue. These neurospheres can be grown up in large quantities in vitro for transplantation into patients.

As mentioned, Reynolds actually developed the neurosphere assay (NSA), which is now widely used to isolate, propagate and enumerate stem cells derived from the CNS. It is now recognised, however, that not all the neurospheres in a culture are derived from stem cells as first thought. About 90 per cent come from progenitor cells. The numbers of stem cells represented by the NSA are largely indeterminate. Reynolds is also developing assays to address this problem.

Proliferation

The second major problem with growing neural stem cells as neurospheres is that only about 10 per cent of them turn into neurons. When the cells are given growth factors in culture to drive proliferation, it seems to push the predominantly down the astrocyte lineage (approximately 90 per cent).

Since generally only one to 10 per cent of transplanted cells survive, the numbers of cells needed for the treatment of one patient becomes unreasonably large.

"People have tried very hard and for a long time and push cells down the neural pathway and basically, it just doesn't work," he says.

Hence, a need existed for a more accurate way of determining and purifying precursor cells. "We have to know what we are transplanting into patients."

Reynolds' team has come up with a new assay, called the neuroblast assay, which increases the number of neurons that are produced from neurospheres. These are then sorted to give a purity of about 90% neurons. The successful implementation of this technology also depends on being able to identify distinct population of cells within the heterogeneous population of stem and progenitor cells.

"We need to know exactly what is in the culture dish, and what each patient receives in a reproducible way."

Variable and indeterminate combinations of neuronal and other CNS cells are the most likely cause for the neurological effects seen with those early transplants into Parkinson's patients. Part of the technology developed by the team at QBI is focused on sorting the expanded cells to address this exact issue.

Ideally, they will be able to take stem cells from an adult donor, grow them up in tissue culture as neurospheres, separate out the neuronal and non-neuronal cell types, and then reproducibly transplant for each individual. The process additionally allows controlled mixing of the sorted cells.

"You may not want to transplant all neurons - you may want to use 60 per cent neurons and 40% astrocytes. Some transplant papers that report better results when neurons are transplanted with astrocytes."

The ultimate aim for this research is to have a renewable and defined source of neural stem cells that can be used for different patients to treat spinal cord injury, stroke, Parkinson's disease and more.

"Obviously this is just the first step, but we now have a way of figuring out what we need to."

[PRINTER FRIENDLY VERSION]

[OTHER STORIES ABOUT UNIVERSITY OF QUEENSLAND, HIS, NSA, ISLAND, CNS, UNIVERSITY OF QUEENSLAND]

[Send Us E-mail](#) | [Privacy Policy](#)

Copyright 2007 IDG Communications. ABN 14 001 592 650. All rights reserved. Reproduction in whole or in part in any form or medium without express permission of IDG Communications is prohibited. Reuters content is copyright 2007 Reuters. [Click for restrictions.](#)

[Ovid.com](#) | [Login](#) | [Technical Support](#)



[Home](#) [About](#) [Products & Services](#) [Contact Us](#)

Acta Neurologica Scandinavica

Current Issue **Archive**

January 2004, 109:1 > Use of gene therapy in central...

ARTICLE LINKS:

[Fulltext](#) | [PDF \(137 K\)](#)

Use of gene therapy in central nervous system repair.

Review article

Acta Neurologica Scandinavica. 109(1):1-8, January 2004.
Tinsley, R.; Eriksson, P.

Abstract:

Recent advances have increased our molecular understanding of the central nervous system disease. In order to realize the clinical benefits of these findings, new molecular models such as CNS gene therapy. Although the field has suffered setbacks, it represents new therapies in the post-genomic world. The development of new vector models of CNS disease, provides evidence suggesting that gene therapy is a viable option. In fact, the first gene therapy clinical trial for Parkinson's disease has been applied in animal models, and how it may be used for other diseases and trauma in human beings. Furthermore, it explores how such augment, more conventional therapeutic approaches.

(C) 2004 Munksgaard International Publishers Ltd.

Search:

GO

[Advanced Search](#)

Ovid News

[Learn](#) about Ovid's new research content and products.

Ovid Events

[Check out](#) where Ovid is across the globe at industry conferences and events worldwide.

Are you affiliated with a leading medical or academic institution?

If so, send your librarian an email recommending titles you've found valuable for research at your institution.

[Ads by Google](#)

Up to 100% Gene Delivery

Order custom lentiviral vectors to stably express your gene or RNAi.
www.Lentigen.com


Copyright © 2005, Ovid. All rights reserved.
Published by Ovid
[Copyright/Disclaimer Notice](#) || [Privacy Policy](#)

[Contact Ovid](#) || [Technical Support](#)

[nature.com homepage](#)

- [Jump to main content](#)
- [Jump to navigation](#)

[Login](#)



- [My account](#)
- [E-alert sign up](#)

- [Subscribe](#)
- [Register](#)

- [Publications A-Z index](#)
- [Browse by subject](#)

Search [Advanced search](#)

What are you looking for?



Journal of
Cerebral Blood
& Metabolism

Perspective

Nature Reviews Neuroscience 7, 75-84 (January 2006) | doi:10.1038/nrn1829

Opinion: Gene therapy: can neural stem cells deliver?

Franz-Josef Müller^{1,2}, Evan Y. Snyder¹ and Jeanne F. Loring¹ [About the authors](#)

Abstract

Neural stem cells are a self-renewing population that generates the neurons and glia of the developing CNS. Neural stem cells have been considered for use in cell replacement therapies in various neurological disorders. An unexpected and potentially valuable characteristic of these cells has recently been revealed — they are attracted to areas of brain pathology such as ischaemic and neoplastic lesions. Here, we speculate that stem cells might be exploited as delivery vehicles for gene therapy in the CNS.


Neural stem cells can be defined operationally as cells that can continuously self-renew and have the capacity to differentiate into all cell types of the CNS.

intermediate and mature cells of both glial and neuronal lineages¹. There are various subpopulations that can be restricted to particular developmental stages or regions of the mature brain, and each of these has specific biological features^{2,3}. It remains unclear whether cultured cells that are derived from this population have the operational definition of neural stem cells — multipotency and the ability to self-renew — are identical to those that have been reported in vivo. In addition, as there are few consensus criteria that can be used to identify neural stem cells, cells known as neural stem cells in one laboratory may differ considerably from similarly named cells in another. For the purposes of this review, we use an inclusive view, assuming that cells that are called neural stem cells by different investigators do have common features that allow generalization. However, we do add the caveat that a particular neural stem cell line or preparation might not apply to all populations (for more details, see Refs 3–6; for reviews, see Refs 2,7,8).

Neural stem cell homing and drug delivery

The migratory abilities of endogenous and exogenous neural stem cells are well known, and it has been hypothesized that these properties, along with the cells' differentiative abilities, might be harnessed for replacement of diseased cells. In 2000, some reports showed for the first time how these cells might be used in a novel way to deliver therapeutic substances to specific sites in the brain^{9,10,11}. These reports showed that cells transplanted into animal models of brain neoplasia were found near metastatic tumour cells far from the site of transplantation^{9,10,11} (Figs 1,2). These observations suggest that neural stem cells engineered to express a specific therapeutic gene might be used to track down and destroy malignant cells. This opens up a possible new realm for neural stem cells: being viewed solely as restorative cell therapeutics, the cells could help to solve one of the most difficult problems in therapy — how to target therapeutic genes to diseased tissues.

Figure 1 | Neural stem cell homing in brain tumours.




a | Transplanted neural stem cells (NSCs) showing tropism for malignant cells in rodent models. Exogenous neural stem cells implanted at sites distant from experimental brain tumours have been shown to migrate towards the tumour. It is believed that this phenomenon could be exploited to track down widespread metastatic CNS pathology and deliver therapeutic systems into brain malignancies. Panels show a time series for murine neural-stem-cell-transfected with the 'luciferase' gene (Luc) and implanted into one hemisphere of experimental animals. Bioluminescence emission imaging of Luc expression for these animals is shown on day 9, day 15, day 22 and day 28 post-implantation. Experimental tumour (black circle in b) was evident from day 15. f–h | Pathotropism of human neural stem cells (NSCs) for malignant cells in rodent models. f | Distribution of the cells (red) within a U87 (a glioblastoma cell line) xenograft (arrows indicate nuclei shown in purple). g | Distant tumour satellite of a U251 (a glioblastoma cell line) xenograft (arrow indicates blood vessel is marked by an asterisk). Note that neural stem cells can migrate transcallosally from one hemisphere to the other and also infiltrate small tumour satellites that have dislodged from the main tumour mass. h | Proximity of NSCs to malignant cells in rodent models.

blood vessel (green) in a U87 xenograft. Panels b–e reprinted, with permission, from Ref. **82** © Panels f–h reprinted, with permission, from Ref. **23** © (2005) Neoplasia.

- **High resolution image and legend (80 KB)**
- **Figures and tables index**
- **Download Power Point slide (290 KB)**

Figure 2 | Determinants of neural stem cell homing to brain tumours for delivery of



Neural stem cells express various receptors for chemoattractant signals as a result of brain path chemoattractants are chemokines such as stromal cell-derived factor 1 (SDF1, also known as CXCL12) and monocyte chemoattractant protein 1 (MCP1, also known as chemokine (C–C motif) chemotactic proteins such as vascular endothelial growth factor (VEGF). Stem cells can be gene enzymes that metabolize non-toxic prodrugs locally, thereby allowing production of the active cytokines that act directly on the tumour or activate immune cells, which, in turn, attack the tumour.

- **High resolution image and legend (116 KB)**
- **Figures and tables index**
- **Download Power Point slide (325 KB)**

The characteristics of neural stem cells make them suitable as therapeutic delivery vehicles for from other cell types that might be considered for this purpose. Unlike fibroblast cells, which migrate from organs, neural stem cells have the potential to integrate seamlessly into the host brain without differentiation. For example, neural stem cells could differentiate into glia or neurons, but are unlikely to become cancerous. Stem cells can be propagated for long periods, and are therefore amenable to the techniques required for gene delivery. Because stem cells can disperse throughout the brain after transplantation, the use of these cells is preferable to multiple stereotactic injections for the delivery of molecules that require distribution. This approach has been used for enzyme replacement in lysosomal storage diseases¹². Another characteristic of neural stem cells for targeted delivery is their tropic behaviour toward neoplasms, which could be exploited to target infiltrating malignant satellites after main tumour resection.

Neural stem cell pathotropism

Neural stem cells (endogenous and transplanted) seem to be attracted to various experimental lesions such as cancers and areas of neurodegeneration. For example, neural stem cells have shown tropism for degenerating spinal cord motor neurons in a transgenic mouse model of **amyotrophic lateral**

1). Neural stem cell cancer tropism is not limited to primary brain malignancies and has also been

Table 1 | Disease models with neural stem cell tropism


- [Full table](#)
- [Figures and tables index](#)
- [Download Power Point slide \(282 KB\)](#)

The fate of neural stem cells in the presence of lesions is not well understood, because most preclinical markers (for example, reduction in tumour burden or survival time after treatment)^{9,16}. In certain lesions, transplanted cells appear to form astrocytes and neurons¹⁷. A glial fate may not be ideal neuronal differentiation that could participate in abnormal, possibly damaging, circuits.

The normal course of neural stem cell development and migration in vivo is controlled primarily in regions of the brain that harbour neural stem cells. The microenvironments surrounding neural stem cells, including astroglia, microglia and endothelial cells, which are important regulators of stem cell generation and maintenance of brain homeostasis^{18,19,20,21,22}. Disturbances in the environment due to disease can affect stem cell behaviour by disrupting the environmental equilibrium and exposing the cells to new factors. For example, gradients of factors such as vascular endothelial growth factor (**VEGF**) and stromal cell derived factor-1 (**SDF1**), which emanate from distant brain lesions, may act as attractants for stem cells^{23,24}.

In attempting to predict the behaviour of stem cells in the brain, it is important to consider both endogenous and transplanted neural stem cells. Endogenous and transplanted neural stem cells are often found in different regions of the brain, but there are some important differences to keep in mind. Cultured neural stem cells expanded in culture well beyond their expected proliferative capacity in vivo. Because culture can alter the phenotype of cells, culture could markedly alter the cells' response to their environment. A recent study showed that exposure to the mitogen epidermal growth factor (**EGF**) may convert neural stem cells to neuronal progenitors⁴. Until more is known about the receptors expressed by neural stem cells and their effects on genetic, epigenetic, transcriptional and translational levels, information about exogenous factors should be used cautiously to interpretations of endogenous stem cell behaviour^{5,6,8}.

The molecular basis of neural cell pathotropism is not well understood and different pathologies can lead to different tropisms. Cultured neural stem cells express a wide variety of receptors that should enable them to respond to factors that emanate from brain pathologies (**Table 2**). Experimental studies show that they home to local sites of pathology after transplantation (**Table 1**). Some factors that can be held responsible for this phenomenon belong to the chemokine family^{25,26} (chemotactic cytokines; **Table 2**). Chemokine and cytokine production is a common feature of brain pathology, stroke and brain malignancy, which suggests that these factors could be important in mediating stem cell tropism^{24,25,26}.


Table 2 | Cytokines potentially involved in neural stem cell pathotropism


Cytokine	Pathotropism
IL-1	+
IL-2	+
IL-3	+
IL-4	+
IL-5	+
IL-6	+
IL-7	+
IL-8	+
IL-9	+
IL-10	+
IL-11	+
IL-12	+
IL-13	+
IL-14	+
IL-15	+
IL-16	+
IL-17	+
IL-18	+
IL-19	+
IL-20	+
IL-21	+
IL-22	+
IL-23	+
IL-24	+
IL-25	+
IL-26	+
IL-27	+
IL-28	+
IL-29	+
IL-30	+
IL-31	+
IL-32	+
IL-33	+
IL-34	+
IL-35	+
IL-36	+
IL-37	+
IL-38	+
IL-39	+
IL-40	+
IL-41	+
IL-42	+
IL-43	+
IL-44	+
IL-45	+
IL-46	+
IL-47	+
IL-48	+
IL-49	+
IL-50	+
IL-51	+
IL-52	+
IL-53	+
IL-54	+
IL-55	+
IL-56	+
IL-57	+
IL-58	+
IL-59	+
IL-60	+
IL-61	+
IL-62	+
IL-63	+
IL-64	+
IL-65	+
IL-66	+
IL-67	+
IL-68	+
IL-69	+
IL-70	+
IL-71	+
IL-72	+
IL-73	+
IL-74	+
IL-75	+
IL-76	+
IL-77	+
IL-78	+
IL-79	+
IL-80	+
IL-81	+
IL-82	+
IL-83	+
IL-84	+
IL-85	+
IL-86	+
IL-87	+
IL-88	+
IL-89	+
IL-90	+
IL-91	+
IL-92	+
IL-93	+
IL-94	+
IL-95	+
IL-96	+
IL-97	+
IL-98	+
IL-99	+
IL-100	+

- [Full table](#)
- [Figures and tables index](#)
- [Download Power Point slide \(248 KB\)](#)

Regulation of neural stem cell tropism

The available information suggests that there are at least three important physiological processes that regulate the behaviour of transplanted neural stem cells: inflammation, reactive astrocytosis and angiogenesis.

Figure 3 | Determinants of neural stem cell pathotropism.

Neural stem cells are attracted by at least three physiological processes that are common to many CNS pathologies: inflammation, reactive astrocytosis and angiogenesis. Pathology-induced CNS inflammation is mediated by activation of microglia and release of cytokines and chemokines, which, in turn, increase the inflammatory reaction (for instance, the interleukin-6, **IL-6**, and monocyte chemoattractant protein 1, MCP1, also known as chemokine (C-C motif) ligand 2, CCL2). Gradients of these factors can also attract neural stem cells. The brain lesion and subsequent inflammation trigger signals emanating from inflammation, activated astrocytes secrete chemotactic factors (for example, SDF1, also known as chemokine (C-X-C motif) ligand 12, CXCL12, and vascular endothelial growth factor, VEGF). These factors can act both as chemoattractants for neural stem cells and as promoters of pathology-induced angiogenesis. Angiogenesis and inflammation-activated endothelial cells enhance neural stem cell homing to the lesion site by secreting chemoattractant factors (such as SDF1), and also offer an atypical, perivascular niche for support and differentiation.

- [High resolution image and legend \(117 KB\)](#)
- [Figures and tables index](#)
- [Download Power Point slide \(326 KB\)](#)


Inflammation. In vitro, microglia can induce neural stem cell migration^{22, 27}. It is an attractive

inflammatory response to brain pathology is the common denominator responsible for the seen cells to disperse brain lesions. The prevailing view, based on studies of multiple sclerosis, epile encephalitis and brain irradiation, is that brain inflammation is detrimental to the CNS in gene ^{29,30,31}. Microglia are the first line in defence against brain pathologies, functioning as a dam; and responding to insults by producing cytokines, which, in turn, initiate further reactive respo cells also release neurotrophins^{29,33}, which would be expected to protect neurons, and microgl can modulate the mobilization of neural stem cells both in vitro and in vivo. This suggests that r initiating and coordinating neural-stem-cell-based brain repair mechanisms^{22, 27, 29, 34}.

Reactive astrogliosis. As inflammatory cytokines are released by microglia in response to a astrogliosis, characterized by hyperplasia, hypertrophy and an increase in glial fibrillary acidic p ³⁶. Triggers and mechanisms of this multifaceted response are not fully understood, but some f proximity of the astrocytes to a CNS pathology, the type of lesion and the types of cytokine prod (**IL-1 β**)^{35, 37}.

Studies of the acute effects of inflammatory signals suggest that certain types of activated astrocyte tissue regeneration and stem cell migration^{38, 39, 40} (**Fig. 4**). For example, reactive astrocytes SDF1, which is at least partially responsible for the attraction of neural stem cells to these lesion involved in the guidance of leukocyte and glial homing toward brain injuries^{41, 42} and can reveal invoked by tissue damage to a phenotype resembling radial glia in the developing brain⁴³. These progenitor migration⁴³. Although some types of glial activation might have beneficial effects, it reactive astrocytes are thought to interfere with neuronal–glial signalling and impede neural pr scars and secreting factors such as slit homologue 2 (**SLIT2**), tumour necrosis factor- α (**TNF α**), hyaluronic acid^{44, 45, 46}.

Figure 4 | Model of activated astrocyte mediation of neural stem cell homing to brain lesions.



CNS injury, hypoxia, microglial activation and the subsequent release of inflammatory cytokine 6 and ciliary neurotrophic factor (CNTF)) invoke complex responses known collectively as gliosis. In response to CNS injury, some mature glia revert to a developmental, radial-glia-like state, and can directly migrate towards brain lesions. Cytokine release also causes transient activation of astrocytes. These transiently activated astrocytes become a source of chemoattractants (such as stromal cell-derived factor 1 (SDF1), vascular endothelial growth factor (VEGF), and chemokine (C-C motif) protein 1 (MCP1)) that act on neural stem cells (NSCs)^{3, 4, 5, 6}. SDF1 may directly

toward brain pathology along non-stereotypical routes⁴. Other factors (such as fibroblast growth factor 1 (IGF1)) supplied by reactive astrocytes support neural stem cell proliferation, but astrocytes proliferate reactively, become hypertrophic and increase their glial fibrillary acidic protein. This eventually results in the formation of a tightly compacted astroglial scar, which is the source of SLIT2, tumour necrosis factor- α (TNF α) and hyaluronan that repel neural stem cells and limit regenerative capacity^{3, 8, 9}.

- [High resolution image and legend \(139 KB\)](#)
- [Figures and tables index](#)
- [Download Power Point slide \(351 KB\)](#)

Angiogenesis. Evidence is emerging for an intimate relationship between CNS morphogenesis and angiogenesis. The basal lamina produced by endothelial cells contains many components that are believed to be in the neurogenic niche^{21, 49}. Therefore, endothelial cells could also be involved in the regulation of brain pathology. Vasculogenesis resulting from brain pathology could enhance neural stem cell mobilization. Chemoattractants such as VEGF. VEGF-mediated homing of stem cells might have a key role in stem cell glioma tropism^{23, 50, 51}. In addition, SDF1 is expressed by endothelial cells as well as pericytes and could be important for attraction of neural stem cells²⁴. Because neural stem cells seem to interact with endothelial cells from the luminal side, adhering and transmigrating in a similar fashion to leukocytes, they can be delivered via the bloodstream. In support of this idea, a recent report shows that neural stem cells in the bloodstream in a mouse model of multiple sclerosis establish atypical niches around blood vessels, maintain an undifferentiated state and appear to suppress the inflammatory process^{53, 54}.

Choosing a vehicle for delivery

There is considerable diversity among neural stem cell lines and they may not all be equally suited for stem cell delivery. A suitable delivery vehicle would be stable in tissue culture and capable of sustained, preferably long-term, release of molecules. The cells should have predictable and appropriate differentiation patterns in culture and survive long term in vivo without forming tumours. For the therapeutic strategy to be effective, the cells should demonstrate responsiveness to the chemotactic signals produced by the type of pathology that they are intended to treat. It would be a means for facile delivery of the cells (for example, via the bloodstream).

Should cell lines be immortalized? Historically, non-tumour cells had to be immortalized sufficiently to facilitate their characterization. Immortalizing cells usually involves introduction of an oncogene to expand beyond the time at which they would normally reach senescence. Identification of genetic markers for culture of non-immortalized neural stem cells has made immortalization less necessary, and the use of immortalized cells as research tools and in clinical settings. Immortalized neural stem cells appear to be atypical of most neural stem cell populations, such as extraordinary migratory abilities in vivo (e.g., crossing the blood-brain barrier) and a higher degree of multipotency, which may increase the probability of tumour formation by

However, there are some cases in which oncogene immortalization is an asset. Safety concerns limit the value of the more pronounced invasiveness and migratory capabilities of immortalized neural stem cells. Immortalization can allow propagation of cells with definable properties almost indefinitely, so that particular traits can be established. Furthermore, if immortalized cells could be shown to be more reliable and much easier to control their quality than the quality of primary cell preparations for use in clinical trials, they could be subjected to much more thorough analysis.

Primary cells for transplantation. Although it can be argued that the ideal cell type for transplant is the endogenous neural stem cell as possible, there are some serious limitations to the use of primary stem cell reconstitution of bone marrow, it is unlikely that a single neural stem cell or small group can regenerate damaged brain tissue, so expansion of cells in culture will be required. How much expansion of a sufficient number of viable stem cells for a successful transplant has to be determined empirically.

In vitro culture creates its own problems. There are many neural stem cell lines and preparations under different conditions from diverse sources and maintained under a wide variety of culture conditions. To be used clinically, an important challenge will be for investigators to agree on a common set of characterization criteria.

Human embryonic stem cell-derived cells: can they be effective and safe? Human embryonic stem (ES) cells have several powerful advantages over other types of stem cell for therapeutic approaches. ES cells are derived from cells of the inner cell mass of blastocyst-stage embryos. Unlike neural stem cells, differentiation of ES cells into all elements of the nervous system. ES cells are also immortal and do not undergo senescence after passage. Perhaps most importantly, the focused efforts to characterize ES cells in many laboratories mean that there are the same well-studied cell populations. This reduces the problems of reproducibility and quality with other stem cells. Human ES cells can be induced to differentiate along neuronal lineages⁵⁹ and they resemble somatic neural stem cells. However, there are several important challenges that arise for their use in therapeutic approaches. First, we have to anticipate that because they have not experienced normal development, ES cells might not develop conventional cellular phenotypes, and this may result in unpredictable behavior. Second, much data are available on the migratory potential of human ES-cell-derived neural stem cell populations compared to primary neural stem cells. Furthermore, populations of ES-derived transplanted cells must be shown to be stable and to characterize undifferentiated populations. It is also important to keep in mind that the use of ES cells raises concerns associated with the derivation of these cells from early embryos.

How will we decide? The decision for or against a certain cell preparation must be based on the balance of benefits and harm, and the modern concept of evidence-based medicine. At present, a widely held view is that stem cell-based approaches will be the most relevant therapeutic systems for targeting brain pathologies. However, there is insufficient evidence for any cell type, and there is a need for comprehensive studies that compare different cell populations. New approaches, such as improved in vivo cell tracking tools, will be important for resolution of these issues.

Neural stem cell-based gene therapy

In the nervous system, replacement of neurons is often considered to be the main goal of cell therapy. However, stem cells, are already being used as gene delivery tools and for rescuing neurons rather than replacing them. One approach to gene therapy is that diseases that are caused by the lack of some crucial protein can be treated by introducing a functional gene using appropriate gene expression vectors. This idea was originally proposed for hereditary diseases such as phenylketonuria. In these diseases, a mutated catabolic enzyme causes a metabolic logjam in the upstream pathway, leading to the accumulation of affected cells and surrounding tissue with accumulating substrates and toxic side products. In the case of gene therapy, the expression of a functional replacement gene in or near the affected area would restore the metabolic pathway. This defined concept has been broadened to include any genetic manipulation of cells or tissue to treat a disease. Gene therapies for **Alzheimer's disease** include targeted expression of choline acetyltransferase, which produces acetylcholine, which therefore results in the localized delivery of small molecules, in this case acetylcholine. The use of stem cells will be used therapeutically depends on the nature of the disease or damage that requires treatment.

Neural stem cells can be genetically transduced in vitro and in vivo^{12, 64}. Currently, the most effective method for introducing genes into neural stem cells is by means of lentiviral vectors; the chief concerns about this approach are the potential for

transgene silencing in situ and that integration of the transgene can activate a nearby oncogene, growing subclones^{58, 65, 66}.

To highlight various stem cell-mediated gene delivery strategies, we discuss in more detail six systems that may benefit from such therapy.


Parkinson's disease. A lack of dopamine in the putamen (caused by the degeneration of innervation) has a central role in the pathogenesis of **Parkinson's disease** (PD). Systemic L-DOPA is an effective treatment for the symptoms of PD early in the course of the disease, but does not prevent degeneration and eventually becomes ineffective. Because the degeneration is relatively localized, cell therapy, and experimental transplants of fetal dopamine neurons into the putamen were performed in patients, the successful transplants seemed to work as dopamine pumps, similar to the systemic treatment, the chief advantage of a transplant being a smoothing of the on–off cycle of symptoms, in which patients are often being unable to move and periods of uncontrollable movement^{67, 68}. A concern about this approach is its variability, which is partly due to the inconsistency of the fetal tissue used for transplant and is a characteristic of the disease in each patient; in a controlled study, the best therapeutic benefit correlated with the best achievable symptomatic improvement using L-DOPA in the same patient⁶⁸. The mechanism is not clear; because of the paucity of functional connections in many transplants, it has been proposed that they were acting more as gene therapy vehicles for dopamine delivery than as replacement neurons. However, they might not be limited to acting as dopamine pumps; in some cases, functional connections have been observed. It has been suggested that the transplanted cells may produce trophic factors that help to protect remaining neurons. Preclinical investigations are testing the use of genetically induced production of neurotrophic factors such as neurotrophic factor (**GDNF**) or VEGF in neural stem cell transplants^{69, 70, 71} (see also the following section on neurotrophic factor delivery in neurodegenerative diseases).

Alzheimer's disease. Alzheimer's disease presents a greater therapeutic challenge than PD, but it is widespread, beginning in the hippocampus, cortex and amygdala, and progressing to other regions of the brain. Strategies for cell and gene therapy are focused on using cells to deliver neurotrophic factors. Neurons are protected from degeneration and to re-activate impaired circuitry in neurodegenerative diseases. A clinical trial using fibroblasts to deliver nerve growth factor (**NGF**) has recently been completed⁷². An adenoviral-associated virus (AAV) to deliver NGF expression vectors directly to the brain.

Most recently, we have proposed that the homing qualities of neural stem cells might be exploited with therapeutic enzymes (F.-J.M. and J.F.L., unpublished observations).

Amyotrophic lateral sclerosis. Experimental studies show that overexpression of growth factors such as growth factor 1 (**IGF1**) or VEGF can have beneficial effects on the course of ALS in animal models. However, this sort of therapy for clinical use is the delivery of these large molecules across the blood–brain barrier. It may be the best means of increasing the production of these factors in situ. Cell therapy might be used to deliver these large proteins to specific areas of the CNS where they can aid in the survival of neurons⁷⁵.

Brain malignancy. Neural stem cells seem to be attracted to certain brain tumours and this allows allowing these cells to be used for local chemotherapy (**Fig. 2**). The main issues under investigation are the optimal choice of stem cell type, and the most effective therapeutic system to use (**Table 3**). So far, several immortalized neural stem cell-like cells in preclinical models. The large variety of therapeutic systems, including viruses, prodrug-converting enzymes, immunomodulatory cytokines, proteins with anti-angiogenic activity and direct anti-tumoural activity^{9, 10, 11, 76, 77}.

Table 3 | Examples of studies on neural stem cell-based gene therapy in animal models

- [Full table](#)
- [Figures and tables index](#)
- [Download Power Point slide \(420 KB\)](#)

Disorders of brain metabolism. Inborn genetic defects affecting the CNS, such as lysosomal storage diseases, are among the most promising targets for stem cell therapy. Dispersion of genetically normal stem cells in the brain can lead to the delivery of missing enzymatic activities^{12, 78}. The current preclinical research objectives for this approach include the timing of treatment, which could be in utero, and the type of stem cell to use. The most straightforward targets are storage diseases that are frequently accompanied by an extremely prominent neuroinflammatory response, such as galactocerebrosidase deficiency (**Krabbe disease** in humans and the twitcher mouse). Significant progress has been made recently in protecting neural stem cells from inflammatory damage and could be applied to this approach.

Neuropathic pain. The delivery of cells and genes to treat certain forms of neuropathic pain is a major goal (Table 3). Potentially therapeutic molecules such as growth factors and neurotransmitters delivered by stem cells could alleviate forms of chronic pain in animal models. An emerging conceptual aspect of these studies is the use of stem cell derivatives — astrocytes and oligodendrocytes) might have another unexpected application; the use of stem cells to alleviate injury-induced central pain by influencing neuronal circuitry and excitability^{80, 81}.

From the pioneering work in PD to the emerging exploration of stem cell therapy for Alzheimer's disease, the enthusiasm for the potential of stem cells for the treatment of various diseases of the nervous system is growing. Gene delivery has the potential to maximize the therapeutic impact of drugs. However, most stem cell-based approaches are still in the preclinical testing phase and will have to pass significant hurdles to become viable therapies.

Summary

Neural stem cells could be exploited as delivery vehicles for therapeutic molecules to treat CNS diseases, moving towards brain pathology, which appears to be mediated at least in part by chemokines. The challenges of this approach are in determining which neural stem cells are appropriate for each application, how they are delivered, and what diseases are suitable targets for this approach.

It is important to remember that the current dominant concept in this field predicts that neural stem cells are primarily for cell replacement therapy. Although experiments continue to be designed with the expectation of yielding surprising new interpretations. We will benefit from remaining receptive to unconventional ideas that will lead us to future discoveries that we cannot imagine today.

Links

DATABASES

OMIM

- **Alzheimer's disease**
- **Amyotrophic lateral sclerosis**
- **Krabbe disease**
- **Parkinson's disease**

Entrez-Gene

- **VEGF**
- **SDF1**
- **EGF**
- **GFAP**
- **IL-1 β**
- **SLIT2**
- **TNF α**
- **GDNF**
- **NGF**
- **IGF1**

FURTHER INFORMATION

- **Clinical trial results, Alzheimer's Disease Education & Referral Center**
- **The official National Institutes of Health resource for stem cell research**

Acknowledgements

We would like to thank N. O. Schmidt for his helpful comments on the manuscript.

Competing interests statement:

The authors declared no competing financial interests.

References

1. Gage, F. H. Mammalian neural stem cells. *Science* 287, 1433–1438 (2000).
 - **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
2. Pevny, L. & Rao, M. S. The stem-cell menagerie. *Trends Neurosci.* 26, 351–359 (2003).
 - **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
3. Soares, S. & Sotelo, C. Adult neural stem cells from the mouse subventricular zone are lim to progenitor cells of similar origin. *Neuroscience* 128, 807–817 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

4. Doetsch, F. , Petreanu, L. , Caille, I. , Garcia-Verdugo, J. M. & Alvarez-Buylla, A. EGF converts precursors in the adult brain into multipotent stem cells. *Neuron* 36, 1021–1034 (2002).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

5. Maslov, A. Y. , Barone, T. A. , Plunkett, R. J. & Pruitt, S. C. Neural stem cell detection, changes in the subventricular zone of mice. *J. Neurosci.* 24, 1726–1733 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

6. Mi, R. *et al.* Immortalized neural stem cells differ from nonimmortalized cortical neurosphere progenitors. *Exp. Neurol.* 194, 301–319 (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

7. Lie, D. C. , Song, H. , Colamarino, S. A. , Ming, G. L. & Gage, F. H. Neurogenesis in the adult nervous system diseases. *Annu. Rev. Pharmacol. Toxicol.* 44, 399–421 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

8. Emsley, J. G. , Mitchell, B. D. , Kempermann, G. & Macklis, J. D. Adult neurogenesis and its progenitors, precursors, and stem cells. *Prog. Neurobiol.* 75, 321–341 (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

9. Benedetti, S. *et al.* Gene therapy of experimental brain tumors using neural progenitor cells.

10. Herrlinger, U. *et al.* Neural precursor cells for delivery of replication-conditional HSV-1 vectors. *Ther.* 1, 347–357 (2000).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

11. Aboody, K. S. *et al.* Neural stem cells display extensive tropism for pathology in adult brain.

gliomas. Proc. Natl Acad. Sci. USA 97, 12846–12851 (2000).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

12. Consiglio, A. *et al.* Robust *in vivo* gene transfer into adult mammalian neural stem cells by Sci. USA 101, 14835–14840 (2004).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

13. Glass, R. *et al.* Glioblastoma-induced attraction of endogenous neural precursor cells is as Neurosci. 25, 2637–2646 (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

14. Chi, L. *et al.* Motor neuron degeneration promotes neural progenitor cell proliferation, in spinal cords of ALS mice. Stem Cells Express 11 Aug 2005 (10.1634/stemcells.2005-0076)

- [Article](#)

15. Brown, A. B. *et al.* Intravascular delivery of neural stem cell lines to target intracranial and non-neural origin. Hum. Gene Ther. 14, 1777–1785 (2003).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

16. Li, S. *et al.* Bystander effect-mediated gene therapy of gliomas using genetically engineered Ther. 12, 600–607 (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

17. Kelly, S. *et al.* Transplanted human fetal neural stem cells survive, migrate, and differentiate Proc. Natl Acad. Sci. USA 101, 11839–11844 (2004).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

18. Song, H. , Stevens, C. F. & Gage, F. H. Astroglia induce neurogenesis from adult neural stem

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

19. Palmer, T. D. , Willhoite, A. R. & Gage, F. H. Vascular niche for adult hippocampal neurog

494 (2000).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

20. Shen, Q. *et al.* Endothelial cells stimulate self-renewal and expand neurogenesis of neural (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

21. Mercier, F. , Kitasako, J. T. & Hatton, G. I. Anatomy of the brain neurogenic zones revisite fibroblast/macrophage network. *J. Comp. Neurol.* 451, 170–188 (2002).

- [Article](#)
- [PubMed](#)
- [ISI](#)

22. Aarum, J. , Sandberg, K. , Haeberlein, S. L. & Persson, M. A. Migration and differentiation directed by microglia. *Proc. Natl Acad. Sci. USA* 100, 15983–15988 (2003).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

23. Schmidt, N. O. *et al.* Brain tumor tropism of transplanted human neural stem cells is indu factor. *Neoplasia* 7, 623–629 (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

24. Imitola, J. *et al.* Directed migration of neural stem cells to sites of CNS injury by the stron chemokine receptor 4 pathway. *Proc. Natl Acad. Sci. USA* 101, 18117–18122 (2004).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

25. Ehtesham, M. *et al.* Glioma tropic neural stem cells consist of astrocytic precursors and th by CXCR4. *Neoplasia* 6, 287–293 (2004).

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

26. Widera, D. *et al.* MCP-1 induces migration of adult neural stem cells. *Eur. J. Cell Biol.* 83,

- [Article](#)
- [PubMed](#)
- [ISI](#)

- **ChemPort**
- 27. Ben-Hur, T. *et al.* Effects of proinflammatory cytokines on the growth, fate, and motility of cells. *Mol. Cell. Neurosci.* 24, 623–631 (2003).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 28. Block, M. L. & Hong, J. S. Microglia and inflammation-mediated neurodegeneration: multiple mechanisms. *Prog. Neurobiol.* 76, 77–98 (2005).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 29. Aharoni, R., Arnon, R. & Eilam, R. Neurogenesis and neuroprotection induced by peripheral inflammation in experimental autoimmune encephalomyelitis. *J. Neurosci.* 25, 8217–8228 (2005).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 30. Ekdahl, C. T., Claassen, J. H., Bonde, S., Kokaia, Z. & Lindvall, O. Inflammation is detrimental to neurogenesis in adult brain. *Proc. Natl Acad. Sci. USA* 100, 13632–13637 (2003).
- **Article**
 - **PubMed**
 - **ChemPort**
- 31. Monje, M. L., Toda, H. & Palmer, T. D. Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 308, 1765–1768 (2003).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 32. Nimmerjahn, A., Kirchhoff, F. & Helmchen, F. Resting microglial cells are highly dynamic in vivo. *Science* 308, 1314–1318 (2005).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 33. Barouch, R. & Schwartz, M. Autoreactive T cells induce neurotrophin production by immune cells in the optic nerve: implications for protective autoimmunity. *FASEB J.* 16, 1304–1306 (2002).
- **PubMed**
 - **ChemPort**
- 34. Wong, G., Goldshmit, Y. & Turnley, A. M. Interferon- γ but not TNF α promotes neuronal outgrowth of murine adult neural stem cells. *Exp. Neurol.* 187, 171–177 (2004).

- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
35. Fernaud-Espinosa, I. , Nieto-Sampedro, M. & Bovolenta, P. Differential activation of microglial cells in the glial scar of the rat optic chiasm. *Glia* 8, 277–291 (1993).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
36. Wang, K. & Walz, W. Unusual topographical pattern of proximal astrogliosis around a cortical lesion. *Neurosci. Res.* 73, 497–506 (2003).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
37. da Cunha, A. *et al.* Control of astrocytosis by interleukin-1 and transforming growth factor β . *Neurosci. Res.* 39–45 (1993).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
38. Spranger, M. *et al.* Regulation of nerve growth factor (NGF) synthesis in the rat central nervous system: the effects of interleukin-1 and various growth factors in astrocyte cultures and *in vivo*. *European Journal of Neuroscience* 15, 100–110 (2003).
- [PubMed](#)
 - [ISI](#)
39. Friedman, W. J. *et al.* Regulation of β -nerve growth factor expression by inflammatory mediators. *Neurosci. Res.* 27, 374–382 (1990).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
40. Liberto, C. M. , Albrecht, P. J. , Herx, L. M. , Yong, V. W. & Levison, S. W. Pro-regenerative microglia in the adult rat brain. *J. Neurochem.* 89, 1092–1100 (2004).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
41. Babcock, A. A. , Kuziel, W. A. , Rivest, S. & Owens, T. Chemokine expression by glial cells in response to injury in the CNS. *J. Neurosci.* 23, 7922–7930 (2003).
- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
42. Wang, K. , Bekar, L. K. , Furber, K. & Walz, W. Vimentin-expressing proximal reactive astrocytes in the rat optic chiasm. *Glia* 42, 100–110 (2004).

rather than proliferation following focal brain injury. *Brain Res.* 1024, 193–202 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

43. Leavitt, B. R. , Hernit-Grant, C. S. & Macklis, J. D. Mature astrocytes transform into trans mouse neocortex that supports directed migration of transplanted immature neurons. *Exp*

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

44. Hagino, S. *et al.* Slit and glypican-1 mRNAs are coexpressed in the reactive astrocytes of th 138 (2003).

- [Article](#)
- [PubMed](#)
- [ISI](#)

45. Fawcett, J. W. & Asher, R. A. The glial scar and central nervous system repair. *Brain Res.* 1

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

46. Back, S. A. *et al.* Hyaluronan accumulates in demyelinated lesions and inhibits oligodendr Nature Med. 11, 966–972 (2005).

- [Article](#)

47. Cleaver, O. & Melton, D. A. Endothelial signaling during development. *Nature Med.* 9, 661

- [Article](#)

48. Carmeliet, P. & Tessier-Lavigne, M. Common mechanisms of nerve and blood vessel wirin

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

49. Alvarez-Buylla, A. & Lim, D. A. For the long run: maintaining germinal niches in the adult

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

50. Bagnard, D. *et al.* Semaphorin 3A–vascular endothelial growth factor-165 balance mediat progenitor cells by the recruitment of shared receptor. *J. Neurosci.* 21, 3332–3341 (2001).

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

51. Zhang, H. , Vutskits, L. , Pepper, M. S. & Kiss, J. Z. VEGF is a chemoattractant for FGF-2-
Cell Biol. 163, 1375–1384 (2003).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
52. Allport, J. R. , Shinde Patil, V. R. & Weissleder, R. Murine neuronal progenitor cells are pr
vasculature via $\alpha 4$ -integrin and SDF-1 α -dependent mechanisms. Cancer Biol. Ther. 3, 838
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
53. Pluchino, S. *et al.* Injection of adult neurospheres induces recovery in a chronic model of i
694 (2003).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
54. Pluchino, S. *et al.* Neurosphere-derived multipotent precursors promote neuroprotection
mechanism. Nature 436, 266–271 (2005).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
55. Ourednik, J. , Ourednik, V. , Lynch, W. P. , Schachner, M. & Snyder, E. Y. Neural stem cell
for rescuing dysfunctional neurons. Nature Biotechnol. 20, 1103–1110 (2002).
 - [Article](#)
56. Ruschenschmidt, C. , Koch, P. G. , Brustle, O. & Beck, H. Functional properties of ES cell-
hippocampus of adult normal and chronically epileptic rats. Epilepsia 46 (Suppl. 5), 174–1
 - [Article](#)
57. Gao, W. Q. & Hatten, M. E. Immortalizing oncogenes subvert the establishment of granule
cerebellum. Development 120, 1059–1070 (1994).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
58. Imren, S. *et al.* High-level β -globin expression and preferred intragenic integration after l
cord blood stem cells. J. Clin. Invest. 114, 953–962 (2004).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
59. Carpenter, M. K. *et al.* Enrichment of neurons and neural precursors from human embryo
383–397 (2001).

- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
60. Tang, Y. *et al.* *In vivo* tracking of neural progenitor cell migration to glioblastomas. Hum.
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
61. Hoehn, M. *et al.* Monitoring of implanted stem cell migration *in vivo*: a highly resolved *in* investigation of experimental stroke in rat. Proc. Natl Acad. Sci. USA 99, 16267–16272 (2002).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
62. Gene Therapy *Wikipedia* [online], <http://en.wikipedia.org/wiki/Gene_therapy>
63. Pizzo, D. P. , Coufal, N. G. , Lortie, M. J. , Gage, F. H. & Thal, L. J. Regulatable acetylcholin cognitive performance. Mol. Ther. 26 Sep 2005 (10.1016/j.mythe.2005.08.001).
- [Article](#)
64. Bharali, D. J. *et al.* Organically modified silica nanoparticles: a nonviral vector for *in vivo* brain. Proc. Natl Acad. Sci. USA 102, 11539–11544 (2005).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
65. Kustikova, O. *et al.* Clonal dominance of hematopoietic stem cells triggered by retroviral g 1174 (2005).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
66. Vroemen, M. , Weidner, N. & Blesch, A. Loss of gene expression in lentivirus- and retrovir cells is correlated to migration and differentiation in the adult spinal cord. Exp. Neurol. 195, 105–115 (2005).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
67. Zurn, A. D. , Tseng, J. & Aebischer, P. Treatment of Parkinson's disease. Symptomatic cell minipumps. Eur. Neurol. 36, 405–408 (1996).
- [PubMed](#)
 - [ChemPort](#)
68. Freed, C. R. , Breeze, R. E. , Fahn, S. & Eidelberg, D. Preoperative response to levodopa is outcome. Ann. Neurol. 55, 896; author reply 896–897 (2004).

- **Article**
 - **PubMed**
 - **ISI**
69. Akerud, P. , Canals, J. M. , Snyder, E. Y. & Arenas, E. Neuroprotection through delivery of factor by neural stem cells in a mouse model of Parkinson's disease. *J. Neurosci.* 21, 8108-8115 (2001).
- **PubMed**
 - **ISI**
 - **ChemPort**
70. Casper, D. *et al.* Enhanced vascularization and survival of neural transplants with *ex vivo* Transplant. 11, 331–349 (2002).
- **PubMed**
 - **ISI**
71. Yasuhara, T. *et al.* Neurorescue effects of VEGF on a rat model of Parkinson's disease. *Bra*
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
72. Tuszynski, M. H. *et al.* A phase 1 clinical trial of nerve growth factor gene therapy for Alzh
- 555 (2005).
- **Article**
73. Kaspar, B. K. , Llado, J. , Sherkat, N. , Rothstein, J. D. & Gage, F. H. Retrograde viral deliv
- mouse ALS model. *Science* 301, 839–842 (2003).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
74. Azzouz, M. *et al.* VEGF delivery with retrogradely transported lentivector prolongs survival
- 429, 413–417 (2004).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
75. Klein, S. M. *et al.* GDNF delivery using human neural progenitor cells in a rat model of AL
- (2005).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
76. Ehteshami, M. *et al.* Induction of glioblastoma apoptosis using neural stem cell-mediated c
- related apoptosis-inducing ligand. *Cancer Res.* 62, 7170–7174 (2002).
- **PubMed**

- **ISI**
 - **ChemPort**
77. Kim, S. K. *et al.* PEX-producing human neural stem cells inhibit tumor growth in a mouse 5965–5970 (2005).
- **PubMed**
 - **ISI**
 - **ChemPort**
78. Snyder, E. Y. , Taylor, R. M. & Wolfe, J. H. Neural progenitor cell engraftment corrects lys VII mouse brain. *Nature* 374, 367–370 (1995).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
79. Monje, M. L. & Palmer, T. Prevention of deficits in neurogenesis with anti-inflammatory a 20040254152 (2004).
80. Hains, B. C. , Johnson, K. M. , Eaton, M. J. , Willis, W. D. & Hulsebosch, C. E. Serotonergi attenuate bilateral hyperexcitability of dorsal horn neurons after spinal hemisection in rat (2003).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
81. Hofstetter, C. P. *et al.* Allodynia limits the usefulness of intraspinal neural stem cell grafts outcome. *Nature Neurosci.* 8, 346–353 (2005).
- **Article**
82. Weissleder, R. & Ntziachristos, V. Shedding light onto live molecular targets. *Nature Med.*
- **Article**
83. Allport, J. R. , Shinde Patil, V. R. & Weissleder, R. Murine neuronal progenitor cells are pr vasculature via α_4 -integrin and SDF-1 α -dependent mechanisms. *Cancer Biol. Ther.* 3, 838-
- **PubMed**
 - **ISI**
 - **ChemPort**
84. Burnstein, R. M. *et al.* Differentiation and migration of long term expanded human neural model of Parkinson's disease. *Int. J. Biochem. Cell Biol.* 36, 702–713 (2004).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
85. Ahn, T. B. , Kim, J. M. , Kwon, K. M. , Lee, S. H. & Jeon, B. S. Survival and migration of tr dopamine cells in the brain of parkinsonian rat. *Int. J. Neurosci.* 114, 575–585 (2004).

- [Article](#)
 - [PubMed](#)
 - [ISI](#)
86. Chen, J. , Magavi, S. S. & Macklis, J. D. Neurogenesis of corticospinal motor neurons extends to adult mice. *Proc. Natl Acad. Sci. USA* 101, 16357–16362 (2004).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
87. Shihabuddin, L. S. *et al.* Intracerebral transplantation of adult mouse neural progenitor cells leads to a marked decrease in lysosomal storage pathology. *J. Neurosci.* 24, 10642–10651 (2004).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
88. Hayashi, T. *et al.* Neural precursor cells division and migration in neonatal rat brain after stroke. *Brain Res.* 1038, 41–49 (2005).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
89. Shear, D. A. *et al.* Neural progenitor cell transplants promote long-term functional recovery after stroke. *Brain Res.* 1026, 11–22 (2004).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
90. Wennersten, A. , Meier, X. , Holmin, S. , Wahlberg, L. & Mathiesen, T. Proliferation, migration and differentiation of neural stem/progenitor cells after transplantation into a rat model of traumatic brain injury. *Brain Res.* 1026, 11–22 (2004).
- [PubMed](#)
 - [ISI](#)
91. Picard-Riera, N. *et al.* Experimental autoimmune encephalomyelitis mobilizes neural progenitor cells from the subventricular zone to undergo oligodendrogenesis in adult mice. *Proc. Natl Acad. Sci. USA* 99, 13211–13216 (2002).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
92. Ben-Hur, T. *et al.* Transplanted multipotential neural precursor cells migrate into the inflamed brain and promote remyelination in experimental autoimmune encephalomyelitis. *Glia* 41, 73–80 (2003).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
93. Totoiu, M. O. , Nistor, G. I. , Lane, T. E. & Keirstead, H. S. Remyelination, axonal sparing,

transplantation of glial-committed progenitor cells into the MHV model of multiple sclerosis (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

94. Ji, J. F. , He, B. P. , Dheen, S. T. & Tay, S. S. Expression of chemokine receptors CXCR4, CXCR5 and CXCR6 in progenitor cells isolated from the subventricular zone of the adult rat brain. *Neurosci. Lett.* 368, 105–108 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

95. Krathwohl, M. D. & Kaiser, J. L. Chemokines promote quiescence and survival of human neural stem cells. *Neuron* 42, 109–118 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

96. Martinez-Serrano, A. & Bjorklund, A. *Ex vivo* nerve growth factor gene transfer to the basal ganglia of middle-aged rats prevents the development of cholinergic neuron atrophy and cognitive impairment. *Acad. Sci. USA* 95, 1858–1863 (1998).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

97. Sun, L. , Lee, J. & Fine, H. A. Neuronally expressed stem cell factor induces neural stem cell self-renewal. *J. Clin. Invest.* 113, 1364–1374 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

98. Jin, K. , Mao, X. O. , Sun, Y. , Xie, L. & Greenberg, D. A. Stem cell factor stimulates neurogenesis in the adult mouse hippocampus. *Invest. Ophthalmol. Vis. Sci.* 43, 311–319 (2002).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

99. Schanzer, A. *et al.* Direct stimulation of adult neural stem cells *in vitro* and neurogenesis *in vivo* by growth factor. *Brain Pathol.* 14, 237–248 (2004).

- [PubMed](#)
- [ISI](#)

100. Martinez-Serrano, A. , Hantzopoulos, P. A. & Bjorklund, A. *Ex vivo* gene transfer of brain-derived neurotrophic factor to intact rat forebrain: neurotrophic effects on cholinergic neurons. *Eur. J. Neurosci.* 8, 727–735 (1996).

- [PubMed](#)
 - [ChemPort](#)
101. Low, W. C. *et al.* Function recovery following neural transplantation of embryonic septal r septohippocampal lesions. *Nature* 300, 260–262 (1982).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
102. Ehtesham, M. *et al.* The use of interleukin 12-secreting neural stem cells for the treatment 62, 5657–5663 (2002).
- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
103. Barresi, V. *et al.* Transplantation of prodrug-converting neural progenitor cells for brain t 10, 396–402 (2003).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
104. Lee, J. *et al.* Cellular and genetic characterization of human adult bone marrow-derived n antiglioma cellular vector. *Cancer Res.* 63, 8877–8889 (2003).
- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
105. Shah, K. *et al.* Glioma therapy and real-time imaging of neural precursor cell migration an 57, 34–41 (2005).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
106. Eaton, M. J. , Santiago, D. I. , Dancausse, H. A. & Whittemore, S. R. Lumbar transplants o neurons alleviate chronic neuropathic pain. *Pain* 72, 59–69 (1997).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
107. Eaton, M. J. , Plunkett, J. A. , Karmally, S. , Martinez, M. A. & Montanez, K. Changes in G the spinal dorsal horn after peripheral nerve injury and promotion of recovery by lumbar t serotonergic precursors. *J. Chem. Neuroanat.* 16, 57–72 (1998).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)

- **ChemPort**

108. Eaton, M. J. *et al.* Transplants of neuronal cells bioengineered to synthesize GABA alleviate pain after sciatic nerve injury. *Transplant.* 8, 87–101 (1999).

- **PubMed**

- **ISI**

- **ChemPort**

109. Eaton, M. J. *et al.* Lumbar transplant of neurons genetically modified to secrete galanin reverses allodynia after partial sciatic nerve injury. *J. Peripher. Nerv. Syst.* 4, 245–257 (1999).

- **PubMed**

- **ISI**

- **ChemPort**

110. Cejas, P. J. *et al.* Lumbar transplant of neurons genetically modified to secrete brain-derived neurotrophic factor reverses allodynia and hyperalgesia after sciatic nerve constriction. *Pain* 86, 195–210 (2000).

- **Article**

- **PubMed**

- **ISI**

- **ChemPort**

111. Lin, C. R. *et al.* Intrathecal spinal progenitor cell transplantation for the treatment of neuropathic pain. *Neurosci. Lett.* 327, 17–24 (2002).

- **PubMed**

- **ISI**

Author affiliations

1. Franz-Josef Müller, Evan Y. Snyder and Jeanne F. Loring are at the Burnham Institute for Medical Research, La Jolla, California, USA.
2. Franz-Josef Müller is also at the Zentrum für Integrative Psychiatrie, Kiel, Germany.

Correspondence to: Franz-Josef Müller^{1,2} Email: fmueeller@burnham.org

- **Previous article**

Top

Nature Reviews Neuroscience

ISSN: 1471-003X

EISSN: 1471-0048

- **About NPG**
- **Contact NPG**
- **Nature jobs.com**
- **Privacy policy**
- **Legal notice**
- **Accessibility statement**

- **RSS web feeds**
 - **Help**
-

© 2007 Nature Publishing Group – partner of AGORA, HINARI, CrossRef and COUNTER

Review

Spinal Cord (2007) **45**, 2–14. doi:10.1038/sj.sc.3101963; published online 1 August 2006

International spinal research trust research strategy. III: A discussion

M Adams¹, T Carlstedt², J Cavanagh¹, R N Lemon³, R McKernan⁴, J V Priestley⁵, G Raisman⁶

1. ¹Research Division, ISRT, Bramley Business Centre, Bramley, Guildford, Surrey, UK
2. ²Royal National Orthopaedic Trust, Stanmore, UK
3. ³The Institute of Neurology, Queen Square, London, UK
4. ⁴Pfizer Limited, Sandwich, Kent, UK
5. ⁵Neuroscience Centre, St Bartholomew's and the Royal London School of Medicine and D
6. ⁶Spinal Repair Unit, Institute of Neurology, Queen Square, London, UK
7. ⁷Netherlands Institute for Brain Research, Amsterdam, The Netherlands

Correspondence: RN Lemon, Sobell Department of Neurophysiology, The Institute of Neurolog
3BG, UK

Abstract

Study design:

Discussion document.

Objectives/methods:

To review the Research Strategy of the International Spinal Research Trust (ISRT), which ident
research that are likely to be beneficial in developing potential treatments for spinal cord injury
intended to both guide the programme of research towards areas of priority and stimulate discu
research. This latest document has been developed to take into account the scientific progress in
previous Research Strategy.

Results/discussion:

The latest scientific developments in research designed to repair the spinal cord and restore fun
might impact on spinal cord injury research are highlighted.

Sponsored by:

ISRT.

Keywords:

spinal cord injury, regeneration, ISRT

Introduction

The International Spinal Research Trust (ISRT) is committed to developing treatments to cure spinal cord injury. For over 25 years it has pursued this goal by funding scientific research proposals from basic to clinical research. Recently, it has recognised the need to improve the scientific basis for assessment of spinal cord injury and committed funding to developing such techniques and training scientists in their use. It is intended that these measures will provide a resource for use in clinical trials worldwide, and will enhance the detection and recovery in such trials.

The record of achievement of the ISRT in promoting basic and clinical research leading to potential cures for spinal cord injury is second to none: it is the pre-eminent UK organisation in this area, and funds international research.

Although by no means the largest funding organisation in this area, ISRT is consistently at the forefront and has developed an enviable reputation for 'punching above its weight' compared with larger organisations. The reasons for this is the ISRT Research Strategy, which has been developed by the Scientific Committee to guide its efforts in particular areas of research.

The first ISRT research strategy document was published in by Harper *et al* in 1996,¹ and was replaced by the second research strategy.² These documents established a coherent research strategy by attracting attention in order to achieve the overall objectives of the ISRT – to repair the damaged spinal cord. The strategy documents identify key areas for funding and support, which enables basic and clinical research proposals, and the Scientific Committee and external reviewers to judge each proposal according to the programme of research to be steered towards areas of priority. In addition, the document has provided a framework for the research programme.

The unprecedented success of spinal cord injury research in the past few years has resulted in significant progress in areas not covered by the previous research strategy. Consequently, ISRT have updated this document to reflect current progress.

The following document, which identifies priorities for basic and clinical research in the coming years, should be read in conjunction with, the existing research strategy.² In general, ISRT expects applications to be influenced by and to refer directly to the themes described in this strategy document, but also to explore novel approaches as they are developed.

In addition to promoting experimental and clinical studies, ISRT considers it vital to promote interaction between scientists and clinicians on the merits, risks and scope for interventions in the aftermath of SCI. This should include the use of inflammatory, anti-proliferative, neuroprotective and immunosuppressive drugs. This should include testing capabilities of existing large study groups worldwide, for example, via the International Consortium for the Study of Paralysis (ICCP) Clinical Trials Workshops (<http://www.campaignforcure.org>), the European Clinical Trials Network (<http://emsci.org>) and the North American Clinical Trials Networks (<http://www.christopherreeve.org/site/c.geIMLP0pGjF/b.1048737/k.322D/NorthAmerica>) with a view to fostering well-founded clinical best practice.

The targets that form the Third ISRT Research Strategy Document reflect current progress in spinal cord injury research.

were highlighted in the earlier Research Strategy, whereas the importance of others has been re previously, overall strategy is divided into two themes: the vertical targets represent experimental capabilities indicate the means by which the vertical targets are likely to be fulfilled.

Vertical targets

- VT1. Early trauma/inflammation and scar tissue
- VT2. Inhibitory and facilitatory influences
- VT3. Guiding regrowth
- VT4. Spared spinal cord cells and fibres
- VT5. Cell- and gene-based therapies
- VT6. Combinatorial therapies
- VT7. Complementary therapies

Horizontal capabilities

- HC1. Animal models
- HC2. Measuring regrowth and restoration of connectivity
- HC3. Clinical trials
- HC4. Collaborative research

Vertical target 1


Minimising the deleterious effects of early trauma, inflammation and scar tissue

Much of the post-traumatic tissue damage and subsequent neurological deficits associated with events that are initiated by the original injury. Thus, spinal injuries comprise a primary zone of secondary injury, and neuroprotective strategies that either reduce or prevent the spread of secondary injury are essential for greater recovery of function.^{3, 4, 5} Many of the mechanisms responsible for these secondary problems are still unknown, and further understanding of the detailed molecular and cellular mechanisms involved in early trauma is vital to provide direction for the rational development of therapeutics to minimise secondary injury following injury. This is the rationale behind treatments such as methylprednisolone,⁶ as GM-1 gangliosides (which also promote plasticity)⁷ and newer developments such as the tetra-

Spinal cord injury should not be regarded in isolation. There are similarities in the mechanisms of ischaemic and traumatic brain injuries,⁹ and some potential therapeutics have already been tested in brain injury. The lessons learned from these (relatively unsuccessful) trials should provide valuable information to the spinal injury community regarding clinical trial issues such as the need for controls and careful dose selection. Therefore, in addition to developing new therapies, an important role of the ISRT is to promote collaboration between basic scientists and clinicians, and the critical evaluation of the merits of these existing and potential treatments.

However, because most neuroprotective strategies have not had particularly substantial effects in clinical trials, it is a priority to increase the understanding of cell death after acute spinal injury and the conditions that lead to it (**Table 1**; VT1.1), in order to develop newer, more powerful therapies.

Table 1 - Vertical target 1: Minimising the deleterious effects of early trauma, inflammation and scar tissue


Full table

It remains important to characterise the effects of injury on major spinal cord components, such as white matter, the composition and effects of the glial scar, vascular effects and the role of inflammatory mechanisms of secondary cell death, and the factors that lead to cyst formation (**Table 1**; VT1.1), the contribution of these events to functional deficits (**Table 1**; VT1.2), and to accurately and model the effects of therapeutic agents on spinal cord function and behaviour. It should then be possible to develop models, both by rational mechanistic drug design and by screening drug libraries in appropriate models, it is also necessary to further our understanding of human spinal injury (**Table 1**; VT1.3), markers of early traumatic injury in humans (**Table 1**; VT1.4).

Exclusions and future issues

Work in non-spinal cord models should be explicitly justified. Funding for the acquisition of new models of inflammation etc. outside the spinal cord, for use in SCI research, will be considered under this Horizontal capability 4.


Vertical target 2

Inhibitory and facilitatory molecules

The inhibitory effects of the CNS on axon growth are well documented, as is the ability of peripheral nerve grafts to promote regeneration. The consensus is that there are several possible reasons why peripheral nerve grafts first is that inhibitory cues that are present in the CNS are absent from the PNS. The second is that the CNS contains neurotrophic factors and contain other regeneration-promoting proteins such as cell-surface proteins that promote inhibition. A third possibility is that inhibitors that are present in PNS grafts are localised and promote regeneration. Thus, it is established that chondroitin sulphate proteoglycans (CSPGs), some myelin proteins present in both the PNS and the CNS, but their expression, stability and localisation/cellular distribution in some types of CNS neurons do not grow into peripheral transplants.¹¹ This may be because of the inhibitory effects of the grafts, or because these CNS neurons are unable to mount an appropriate response of regeneration.

Therefore, research into inhibitory and facilitatory molecules falls into three categories: the discovery of mechanisms of inhibition and facilitation of axon growth (**Table 2**; VT2.1), the production of reagents that either increase inhibition or increase facilitation (**Table 2**; VT2.2), and the investigation of how other approaches, such as cell transplantation, interact with inhibitory/facilitatory molecular mechanisms (**Table 2**; VT2.3).

Table 2 - Vertical Target 2: Research into inhibitory and facilitatory molecules.


Full table

The idea that the presence of inhibitory molecules causes regeneration failure was proposed by culture¹² and then in the injured spinal cord,¹³ on the basis of a molecule present on the surface. This molecule is now known as Nogo. Blocking Nogo promotes either regeneration¹³ or beneficial sp

Nogo is one of several inhibitory molecules that are associated with CNS myelin, with other can glycoprotein and oligodendrocyte myelin glycoprotein. Subsequently, inhibitory molecules of th families have been found to be associated with astrocytes and fibroblasts^{16, 17, 18, 19, 20} and a c these various inhibitory molecules has been described.²¹ It is likely that increased molecular ch inhibitory pathway will help in the development of new therapies.

In addition to the lack of inhibitory molecules, it is proposed that peripheral nerve grafts support secretion of trophic factors. Many studies have shown that growth factors upregulate RAGs and growth,^{22, 23, 24} and genetically modifying transplants to overexpress neurotrophic factors might. Thus, an effective combination that enhances positive factors and reduces negative ones is an at

So far the molecular response to neural injury has been studied mainly at the level of changes in proteins. This has led to the discovery of several molecules that are important in regeneration. 7 projects have identified most human and rodent genes. This enables high-throughput screening proteins, which will be instrumental for elucidating the molecular mechanisms that underlie reg in a more complete picture of the molecular changes that occur in neurons and glial cells after in of proteomic techniques coupled to web-based databases and data-analysis tools is likely to ider to pinpoint novel targets for pharmacological and cell- and gene-based intervention strategies. (processes that underlie regeneration is still very limited, so incorporating genomic and proteom neuroregeneration research is vital to progress in this field.

Vertical target 3

Guiding regrowth and establishing appropriate connections

Several existing therapies promote the regeneration of injured axons in long, white-matter path being developed to bridge spinal injury sites using synthetic biomaterial implants.^{26, 27} However, special environment that these therapies provide, to cross scar tissue associated with the injury beyond the scar has proved a major problem.


Given that the number of axons that do regenerate and regrow is likely to be small compared with important to understand how fibres behave once they have reconnected with the spinal cord beyond extreme, it might not be necessary to guide regrowing fibres to their appropriate targets if intrinsic CNS can exploit any connectivity to achieve some recovery of function. At the other extreme, new ineffective or even add to secondary problems such as spasticity and increased pain levels. We are factors that guide axons to their eventual target by attraction and repulsion, including factors th

tracts until they descend (or ascend) to the level appropriate to penetrate the grey matter.^{28, 29} matter, ephrins, netrins, semaphorins and related molecules guide axons and regulate midline c termination.

It is probable that the main national research funding bodies will continue to invest in this area expanding the 'library' of agents that guide growing and regrowing axons. ISRT should adopt a knowledge into spinal injury studies and promote research into:

- The nature and temporal course of new synaptic connectivity after SCI, using histological (Table 3 ; VT3.2).
- The key guidance/trophic factors – how they are affected by SCI, particularly in regions at (Table 3; VT3.1, VT3.2), and whether their expression can be modified to encourage app VT3.3).
- The potential of axonal sprouting and synaptic plasticity for regeneration and useful re-in
- The ability of trophic factors to restore axonal growth *per se* and to influence the re-establ (Table 3; VT3.4).

Table 3 - Vertical Target 3: Guiding regrowth and establishing appropriate connec



[Full table](#)

Vertical target 4

Assessing the natural history of SCI and optimising spared spinal cord cells and ne

Except for a few open or penetrating injuries, there is spared spinal cord tissue in most cases of complete. Tissue sparing has important consequences, both in terms of the function of the resid plasticity. In animal models, fibres passing through the ventrolateral funiculi, including reticul important for functional recovery of hindlimb locomotor function.³¹ In humans, the importance clear, while damage to corticospinal fibres has more serious effects.³²

The minimal sparing of white matter, in terms of either area or axonal number, that is compatil cats,³³ <25% for non-human primates³⁴ and <10% for humans.³⁵ Minimal deliberate moveme dorsal and plantar flexion of the foot, has been reported in humans with only 3.5–10% of cortic level.³⁵ The idea that a limited number of nerve fibres is sufficient for function has encouraged formulated in the belief that a few new axons that cross the lesion site and connect somewhere v recovery.

After total spinal cord transection in laboratory animals, weight-supported, unassisted stepping circuitry below the level of the transection, without supraspinal control.¹¹ This is not the case in motoneurons pools can be activated by proprioceptive sensory inputs generated by treadmill e cases of incomplete SCI.³⁶ The surviving supraspinal motor input to the spinal cord is insufficie


muscles of a single joint or to simultaneously inhibit antagonist muscles, which indicates that v partially related to function.³⁷ After complete SCI, it appears that local, segmental, propriocept generate patterned muscle activity, but not in a sustained manner.^{36, 38}

In animals and humans, plasticity in the motor systems has been shown at different levels from Erroneous connections made after an injury persist for many years.^{40, 41} Early after the injury t that might be occupied by either inappropriate supraspinal tract axons or local interneurons. Th the development and establishment of aberrant reflexes that might be counterproductive in eith function, for example, spasticity or aberrant reflexes. In addition, the occupation of these sites t might prevent appropriate regenerating fibre systems reconnecting to the right circuits. One wa specific neuromodulatory treatments. The development of techniques that maintain the neuron as far as possible within its 'normal' state, and thereby prepare this tissue for successful interve

Plasticity in the sensory systems, such as collateral sprouting, is well known and might account of sensation below the level of the lesion and the development of neuropathic pain. Imaging stu sensation after SCI is accompanied by reorganisation of the somato-sensory cortex.^{42, 43} Howe explain referred sensation from the viscera perceived in deafferented areas. Nevertheless, larg other CNS sites such as thalamus, brain stem, cuneate nucleus and spinal cord.^{44, 45, 46, 47} Thi pre-existing connections⁴⁸ and actual axonal sprouting.⁴⁹ To achieve functional recovery of sei considered include manipulating the biological environment to promote regeneration and funct strategies that increase reorganisation in the CNS.

It is important to define the structure and function of the remaining spinal cord tissue (**Table 4** enhance its functional capacity most effectively (**Table 4**; VT4.2, VT4.3). The clinical consequ and below the lesion site compared with damage to long fibre systems remains to be establish treatments that are designed to replace lost neurons are to be considered alongside those that le Even if nonfunctional, remaining tissue might have a valuable role in the effects of future interv as a scaffold for new growth.

Table 4 - Vertical target 4: Assessing the natural history of SCI and optimising spa fibres.



Full table

Vertical target 5

Cell- and gene-based therapies

Since the last strategy review in 2000,² major progress has been made in the areas of cell and g includes grafting with fully differentiated tissue such as peripheral nerves, inflammatory system macrophages, CNS-resident cells such as oligodendrocytes and olfactory ensheathing cells, and Stem cells are attractive theoretically because it is envisioned that they will respond to cues and

organ systems, such as the foetal liver following injury and adult heart tissue after cardiac infarct. Embryonic stem cells are less appropriate because they can develop into cancerous teratomas. Tissue of both foetal and adult neural origin are preferred currently. Gene-based therapies where a therapeutic gene is expressed directly in the injured spinal cord or neural transplants are gene transplantation.

Cell-based therapies

Cell-based therapies might act in several, distinct ways: (i) as a potential source of either trophic factors to improve the function of pre-existing spinal cord neurons; (ii) transplanted stem cells might develop into neurons and remyelinate regenerating axons; (iii) transplanted stem cells might develop into functional spinal cord tissue to replace those damaged by injury; and (iv) transplanted cells can serve as a substrate to support axonal growth. The mechanisms predominate in studies that have reported restoration of spinal cord function and the different cell types act in different ways. The use of cells as a source of multiple trophic factors to provide the environment for neuronal regeneration impacts on Vertical targets 2–4.

There has been some work to identify the cell types that are generated *in vivo* after stem cell transplantation. Undifferentiated, neurospheres in culture can generate all types of neural cell. However, although they survive when grafted into the rat spinal cord, they only differentiate into astro- and oligodendrocytes. The adult spinal cord provides the molecular cues for glial, but not neuronal, differentiation.⁵⁰ Current procedures for differentiating, isolating and transplanting them need to be perfected. Disappointingly, transplanting neural stem cells leads to an increase in pain levels (allodynia), which is associated with the spinal cord. However, forcing stem cells into a distinct lineage before transplantation avoids this outcome.⁵² The benefits of grafting differentiated, purified cells require further study. These data suggest that preclinical studies should specifically examine the adverse effects of cell therapies.

Cell-based therapies require that the cells are readily obtainable, easy to expand and bank, and that they provide sufficient and appropriate axonal repair. Until large-scale, well-characterised adult and differentiated cells are available, bone marrow mesenchymal stem cells (MSCs) are an attractive source that allows autologous transplantation where the subject receives their own bone marrow. Transplanted unpurified MSCs improve remyelination after spinal cord injury, and several studies achieve modest functional recovery.^{54, 55, 56} Differentiated Schwann cells might further improve the outcome,⁵⁷ as might selection of MSCs from differentiated cells. Different cell types produce similarly effective cells, presumably because of the repertoire of cytokines and modulators.

An alternative source of adult transplantable cells with repair potential are olfactory ensheathing cells generated from cultures of primary olfactory tissue: although both are essential for the reparative relationship of the two cell types is not fully understood.⁶⁰ Olfactory ensheathing cells encourage axonal growth. However, they are likely to be more effective when combined with other treatments (see vertical targets). Inhibitory cues of the scar tissue with chondroitinase ABC and providing a Schwann cell bridge. Olfactory ensheathing cells promotes greater functional recovery in a rat model than these treatments and

Gene therapy

The first viral vectors used to express a therapeutic gene in the nervous system were imperfect b response. These problems inspired the development of improved 'minimal' vectors based on ad These viral vectors carry a transgene under a strong viral or cellular promotor, but are virtually Adeno-associated viral-vector-mediated expression of neurotrophins has been successful in resc root avulsion and reversing atrophy of chronically lesioned rubrospinal neurons.^{62, 63} In additi gene transfer, cellular transplants have been genetically modified *ex vivo* before transplantatio guides. The steady advances made in combining new viral vector systems with a range of promi holds fascinating perspectives for the development of new spinal cord repair strategies (reviewe

Although there has been much progress in the area of cell therapy, significant questions remain in gene therapy are: (i) how to enhance the level of expression of the transgene; (ii) control of th (iii) the difficulty in predicting and controlling the cell types that are transduced, and some cells then others (eg scar tissue can hardly be transduced for, as yet, unknown reasons); and (iv) the and dominant-negative proteins to overcome local action of inhibitory proteins is in its infancy. powerful technique that might be used to overcome inhibition and to enhance the expression of

Table 5 - Vertical target 5: Cell- and gene-based therapies.

Full table

Vertical target 6

Combining therapies

Several independent mechanisms contribute to the outcome of SCI. Therefore, it seems reasona are directed at one specific injury mechanism are likely to have limited overall efficacy, and that might achieve a greater benefit and increase recovery.⁶¹ Some published studies have combined plasticity-promoting drugs to provide proof-of-principle of this concept. It is likely that more eff and combination therapy is likely to be a cornerstone of future strategies following SCI. Howeve between different interactions, interpreting the effect of combining potential treatments require

The potential complementarity of different therapies is crucial, and funding will only be conside cogent case for combining individual approaches (**Table 6**).

Table 6 - Vertical Target 6: Combining therapies.

Full table

Vertical target 7


The complementary role of different forms of electrotherapy for recovery of function after SCI

The aims of ISRT is to develop a long-term, effective treatment for SCI, based on better understanding neurobiological mechanisms of injury and repair. Originally, therefore, functional electrical stimulation, electrotherapy and intensive physiotherapy were not considered central to the research strategy. However, involve actual repair of the injury. However, in the past few years ISRT and the SCI community position.

The primary purpose of FES is to activate paralysed groups of muscles; for example, FES implants for control and to assist standing, locomotion and hand grasp. However, it is clear that, in addition, FES also has long-term 'secondary' effects on central sensorimotor mechanisms^{65, 66} that affect plasticity of surviving fibres to make a greater contribution (see Vertical target 4).

A related point is that the development of the normal spinal cord and, probably, regeneration of at least in part, activity-dependent processes; electrotherapy methods can be used to promote neuroplasticity after injury, which cannot be generated voluntarily by the patient. Finally, we know that activity in the brain above the level of the spinal injury, including the cerebral motor areas, cerebellum and basal ganglia, can be affected by SCI and show different patterns during both attempted and imagined movements. Thus, FES, and other forms of electrotherapy might all complement other, more invasive therapies and boost therapeutic effectiveness (Table 7).

Table 7 - Vertical target 7: The complementary role of different forms of electrotherapy for recovery of function after SCI.



Full table

Some forms of FES are invasive (eg sacral or lumbar root stimulators and intraspinal microstimulation). Some individuals refuse such implants because they expect a more permanent cure to be developed in the future, or because of the chances of inclusion in future trials and treatment.

Given that the principal aim of ISRT is a long-term treatment that provides effective repair of SCI, what are the development of FES and other electrotherapeutic approaches? ISRT should promote research in


- Approaches that demonstrate the complementary role of FES in improving outcome of other treatments for activity-dependent plasticity.
- Studies to determine the extent to which defined FES paradigms improve real-world tasks in individuals with SCI.
- The use of noninvasive methods that offer clear prospects of functional recovery, especially central activity-dependent plasticity.
- Improved outcome measures to assess functional improvements provoked by noninvasive methods.
- The development of long-term, stable, electrotherapy techniques that complement other treatments.

Horizontal capability 1

Animal models

Effective experimental models are crucial for understanding the basic biology of SCI and developing treatments (Table 8). Two common approaches are to use (i) a model that aims to mimic as closely as possible clinically (ie contusion injuries), and (ii) a model in which specific tracts or pathways are lesioned to study a particular system to injury and its capacity for regeneration. Both approaches have their benefits and their limitations. Treatments should be evaluated in both before they are developed for use in humans. Transected animal injury model, whereas contusion represents the typical injury mechanism in humans. A contusion model, which greatly exceeds the extent of neuronal damage in the transected spinal cord. At the level of the lesion, loss of alpha-motoneurons and roots associated with spinal cord contusion is little addressed in the current literature (Vertical target 4), it has direct implications for rehabilitation strategies and functional outcome. The current literature for degradation of neuronal function below the level of lesion in chronic, complete SCI.³⁸ The need for a regeneration-inducing therapy needs to be evaluated. In addition, the prerequisites to facilitate regenerating tract fibres and to maintain neuronal function in the postacute stage have still to be determined.

Table 8 - Horizontal capability 1: Animal models.



Full table

Although the majority of SCI studies to date have involved rats, genomic approaches are carried out in mice. The development of a mouse model of SCI had priority in the 2nd ISRT strategy document.² Now, the use of knockout and transgenic mice is likely to provide insights into the molecular components of SCI.

In each laboratory species used, the requirements of an animal model are that it is quantitative, permanent records that are open and available to other researchers. The results obtained with a model should be reproducible when used independently by other research teams.

Ideally, experiments should have a sequential design that includes:

- Longitudinal observation of the behaviour in normal animals to establish the level of variability and to stabilise learning curves.
- Longitudinal observations of the same parameters after lesion to establish the degree of variability and the natural history and evolution of postlesional changes that occur without any intervention.
- The therapeutic intervention should be applied only when the postlesional situation is stable and a baseline assessment carried out as above.
- Variation must be related to the normal population variation. Postlesional variation should be related to the normal population variation because animals cannot be assumed to be uniform, and correlation with the lesion histology and location of the lesions that are associated with specific effects. In addition, posttherapeutic assessment of individual variation in histological parameters of recovery (eg number of fibres regenerating) should give valuable additional information.

Horizontal capability 2


Developing methods to measure regrowth and restoration of function

The ability to determine accurately the extent of anatomical regeneration, physiological connect crucial for identifying a successful repair strategy. The current lack of reliable assessments of th spontaneous and treatment-induced recovery in laboratory animals and humans are fundament and potential clinical trials for SCI (see Vertical target 4).

The American Spinal Injuries Association (ASIA) score is not always fully reliable. For example, workers^{67, 68} demonstrate that some patients who are classified as ASIA A (complete SCI) can EMG with pudendo-anal reflex (PAR). This sacral reflex is modulated by either voluntary or sup measure of spinal cord injury.⁶⁹ Measuring improvements in the PAR and similar systems migl progressive, postinjury changes in function, and outcome measures for future interventional tre

Methods to detect the partial preservation of long fibre tracts are also needed because present t unable to detect subtle neurological improvements. Assessments that reflect the whole clinical p spasticity, are also needed (**Table 9**). Although spasticity is useful for some abilities such as tra functional outcome for the patient.

Table 9 - Horizontal capability 2: Developing methods for measuring regrowth and



Full table

To some extent neurophysiological and functional assessments can differentiate between the co compensation, neuronal plasticity and regeneration to improvement of function following SCI.⁷ in particular, should give information about the impact of any new interventional therapy on th and of the peripheral nervous system.

Most patients recover significant function without intervention, and deterioration of SCI patient uncommon.⁷¹ For example, most quadriplegic patients recover one spinal level of motor functio SCI, it is difficult to predict the preservation of discrete longitudinal fibre tracts and the likeliho of sensory and motor function. Difficulty in distinguishing between post-treatment improvemen might occur without intervention creates potential problems for interpreting the results of clinic new techniques are needed to assess more effectively spinal cord tissue that is spared after clinic and to predict accurately any spontaneous recovery of function³⁷ (**Table 9**). Collection at multi physiological data mapping the natural history of changes in function in the period immediately

Imaging the site of injury, for example, by MRI can indicate continuity across a lesion but does not functional. Currently, electrophysiological assessments of sensory and motor tract function (eg : indicate the presence of large, myelinated fibres in the dorsal columns but not finer fibres in, for and recovering or remyelinating fibres. Therefore, ways to identify different fibre tracts are need electrodes, which recognise unique patterns of activity in discrete fibre tracts. Functional imagin


developed, and it is likely that the combination of functional imaging with selective stimulation pathways through, for example, contact-heat-evoked-potential stimulation of C and A delta fibre analysis of baseline and functional improvements from interventions.

Horizontal capability 3

Clinical trials

The many issues that surround optimisation of Clinical Trials of SCI treatments were discussed Clinical Trials Workshop on SCI.⁷³ Many SCI and other relevant (eg regulatory) communities w standards and guidelines for valid clinical trials could be developed and broadly accepted. One c establishment of a working group to bring forward detailed guidelines on how to develop clinica effective manner. Clearly, several of the issues relate to the adequacy of the animal model that is launch a clinical trial⁷⁴ (**Table 10** and Horizontal capability 1).

Table 10 - Horizontal capability 3: Clinical trials.



Full table


As a more general consideration, Research Governance applies to all who fund research proposals, research, and host research in their organisation. This is the process that sets standards and defines these, requires monitoring and assessment, and improves research quality and safeguards the public scientific quality, promoting good practice, and preventing poor performance and misconduct. At the highest clinical standards, ISRT must ensure that its governance structures are appropriate to a Strategy.

Horizontal capability 4

Promoting collaborative research

The complexity of SCI in humans is such that multidisciplinary approaches are needed to understand therapies to treat it. It is generally accepted that there is a need to continue to improve communication involved in spinal cord injury research. Therefore, ISRT regards collaboration as one of the keys. To this end, it encourages collaborations between the researchers it funds, both clinical and basic.

Table 11 - Horizontal capability 4: Promoting collaborative research.



Full table

It is important for scientists to understand the general and specific problems associated with SCI. It might be necessary to foster the training and career progression of a new 'breed' of clinical scientists for clinical trials in SCI. Such individuals should be familiar with basic science and have clinical knowledge of the design, execution and evaluation of clinical trials. They should also be able to evaluate research of SCI patients and other clinicians. This is important because what might appear to be an exciting idea might have limited potential for translation to the clinic because of gaps in understanding between basic and clinical scientists and clinicians.

A single centre where different experts, such as basic and clinical scientists and clinicians who work together, is probably most effective. From such a hub, a spoke organisation should be established to exchange ideas and compounds with other units, and to coordinate with other centres to enable sufficient patients to be recruited in time.

In addition, collaboration between basic researchers and clinicians should help to evaluate the mechanisms of complete SCI and, consequently, to better understand neuronal plasticity and degradation. This involves understanding the different factors in determining the severity of functional loss after SCI, such as demyelination site, and link them to therapeutic approaches. An example might be the maintenance of neuronal function in specific, early-onset, functional training.^{75, 76}

Contacts or collaborations outside SCI research should also be encouraged to make use of existing knowledge and treatments and to ensure that mistakes are not repeated. For example, knowledge on the use of stem cells is more advanced in other areas of research such as haematology and cardiology, and this should be shared.

Collaborations with industry should be encouraged, both for support and as a source of new drugs, as a means of promoting international meetings where a wide spectrum of different aspects of SCI research can be discussed.

Conclusions

This latest Research Strategy from the ISRT builds on the previously published strategies^{1, 2} by incorporating recent advances in basic and clinical research that are relevant to restoration of function following SCI. By experience, identifying individual themes of basic and clinical research enables ISRT to focus resources on areas that would benefit from particular attention, and targeting specific research areas in this way maximises the effects of the available funding. As a research-based charity, ISRT intends to grant funding directly by the themes described in this strategy document. However, this is not to say that other areas should be considered should there be sufficiently strong evidence of their potential.

In keeping with the policy of promoting debate between all interested parties, another purpose of this Strategy is to stimulate discussion of the relative merits of the themes and approaches that are considered. This is deliberately wide-ranging and inclusive with respect to the themes described, and individual views on these approaches are likely to differ. By promoting this discussion, ISRT hopes to encourage dialogue between patients and other interest groups about the many issues that are involved in developing and validating new therapeutic advances in the near future.

References

1. Harper GP, Banyard PJ, Sharpe PC. The International Spinal Research Trust's strategic approach to developing treatments for the repair of spinal cord injury. *Spinal Cord* 1996; 34: 449–459. | [PubMed](#)
2. Ramer MS, Harper GP, Bradbury EJ. Progress in spinal cord research: a refined strategy for the future.

- trust. *Spinal Cord* 2000; 38: 449–472. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
3. McDonald JW, Sadowsky C. Spinal-cord injury. *Lancet* 2002; 359: 417–425. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
4. Profyris C, Cheema SS, Zang D, Azari MF, Boyle K, Petratos S. Degenerative and regenerative cord injury. *Neurobiol Dis* 2004; 15: 415–436. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
5. Ramer LM, Ramer MS, Steeves JD. Setting the stage for functional repair of spinal cord in *Cord* 2005; 43: 134–161. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
6. Bracken MB *et al.* Methylprednisolone or tirilazad mesylate administration after acute spinal cord injury. Results of the third National Acute Spinal Cord Injury randomized controlled trial. *J Neurosurg* 2000; 92: 706–717. | [PubMed](#) | [ISI](#) | [ChemPort](#) |
7. Geisler FH, Coleman WP, Grieco G, Poonian D. The Sygen multicenter acute spinal cord injury study. *Spinal Cord* 1998; 36: 115–124. | [Article](#) | [PubMed](#) | [ChemPort](#) |
8. Stirling DP *et al.* Minocycline treatment reduces delayed oligodendrocyte death, attenuates inflammation, and improves functional outcome after spinal cord injury. *J Neurosci* 2004; 24: 2182–2190. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
9. Leker RR, Shohami E. Cerebral ischemia and trauma-different etiologies yet similar mechanisms. *Brain Res Brain Res Rev* 2002; 39: 55–73. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
10. Wahlgren NG, Ahmed N. Neuroprotection in cerebral ischaemia: facts and fancies – the new paradigm. *Cerebrovasc Dis* 2004; 17: 153–166. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
11. Anderson PN, Campbell G, Zhang Y, Lieberman AR. Cellular and molecular correlates of the transition of mammalian CNS axons into peripheral nerve grafts. *Prog Brain Res* 1998; 117: 211–232. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
12. Schwab ME, Thoenen H. Dissociated neurons regenerate into sciatic but not optic nerve after transplantation. *J Neurosci* 1985; 5: 2415–2423. | [PubMed](#) | [ISI](#) | [ChemPort](#) |
13. Schnell L, Schwab ME. Axonal regeneration in the rat spinal cord produced by an antibody that neutralizes growth inhibitors. *Nature* 1990; 343: 269–272. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
14. Chen MS *et al.* Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature* 2000; 403: 434–439. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
15. Bareyre FM, Kerschensteiner M, Raineteau O, Mettenleiter TC, Weinmann O, Schwab ME. A new intraspinal circuit in adult rats. *Nat Neurosci* 2004; 7: 269–277. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
16. McKeon RJ, Höke A, Silver J. Injury-induced proteoglycans inhibit the potential for laminar astrocytic scars. *Exp Neurol* 1995; 136: 32–43. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
17. Bradbury EJ *et al.* Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature* 2002; 416: 636–640. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
18. Bundesen LQ, Scheel TA, Bregman BS, Kromer LF. Ephrin-B2 and EphB2 regulation of axon-astrocyte interactions in response to spinal cord lesions in adult rats. *J Neurosci* 2003; 23: 7789–7800. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
19. Willson CA, Miranda JD, Foster RD, Onifer SM, Whittemore SR. Transection of the adult spinal cord and ligand expression. *Cell Transplant* 2003; 12: 279–290. | [PubMed](#) | [ISI](#) | [ChemPort](#) |
20. Pasterkamp RJ *et al.* Expression of the gene encoding the chemorepellent semaphorin III in the component of neural scar tissue formed following injuries of adult but not neonatal CNS. *J Neurosci* 2000; 20: 166–177. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
21. Filbin MT. Myelin-associated inhibitors of axonal regeneration in the adult mammalian CNS. *Neuron* 2001; 30: 713–724. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
22. Grill R, Murai K, Blesch A, Gage FH, Tuszynski MH. Cellular delivery of neurotrophin-3 promotes axonal growth and partial functional recovery after spinal cord injury. *J Neurosci* 1997; 17: 5560–5572. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
23. Ramer MS, Priestley JV, McMahon SB. Functional regeneration of sensory axons into the spinal cord. *Nature* 2000; 403: 312–316. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
24. Lu P, Jones LL, Snyder EY, Tuszynski MH. Neural stem cells constitutively secrete neurotrophic factors and support extensive host axonal growth after spinal cord injury. *Exp Neurol* 2003; 181: 115–129. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
25. Bareyre FM, Schwab ME. Inflammation, degeneration and regeneration in the injured spinal cord. *Trends Neurosci* 2003; 26: 555–563. | [Article](#) | [PubMed](#) | [ChemPort](#) |

26. Geller HM, Fawcett JW. Building a bridge: engineering spinal cord repair. *Exp Neurol* 2001; 136. | [Article](#) | [PubMed](#) |
27. King VR, Henseler M, Brown RA, Priestley JV. Mats made from fibronectin support orient spinal cord of the adult rat. *Exp Neurol* 2003; 182: 383–398. | [Article](#) | [PubMed](#) | [ChemPort](#) |
28. de Wit J, Verhaagen J. Role of semaphorins in the adult nervous system. *Prog Neurobiol* 2002; 66: 267. | [Article](#) | [PubMed](#) | [ChemPort](#) |
29. Gallo G, Letourneau PC. Regulation of growth cone actin filaments by guidance cues. *J Neurosci* 2002; 22: 102. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
30. Skaper SD, Moore SE, Walsh FS. Cell signalling cascades regulating neuronal growth-promoting factors. *Neurobiol* 2001; 65: 593–608. | [Article](#) | [PubMed](#) | [ChemPort](#) |
31. Noga BR, Kriellaars DJ, Jordan LM. The effect of selective brainstem or spinal cord lesion stimulation of the mesencephalic or pontomedullary locomotor regions. *J Neurosci* 1991; 11: 1700. | [PubMed](#) | [ChemPort](#) |
32. Nathan PW. Effects on movement of surgical incisions into the human spinal cord. *Brain* 1977; 100: 102. | [Article](#) | [PubMed](#) | [ChemPort](#) |
33. Blight AR. Cellular morphology of chronic spinal cord injury in the cat: analysis of myelin. *Neuroscience* 1983; 10: 521–543. | [Article](#) | [PubMed](#) | [ChemPort](#) |
34. Eidelberg E, Story JL, Walden JG, Meyer BL. Anatomical correlates of return of locomotor function after spinal cord injury in cats. *Exp Brain Res* 1981; 42: 81–88. | [Article](#) | [PubMed](#) | [ChemPort](#) |
35. Hofstetter CP *et al.* A review of the neuropathology of human spinal cord injury with emphasis on the cervical region. *Cord Med* 1999; 22: 119–124. | [PubMed](#) | [ChemPort](#) |
36. Maegele M, Muller S, Wernig A, Edgerton VR, Harkema SJ. Recruitment of spinal motor neurons after stepping after human spinal cord injury. *J Neurotrauma* 2002; 19: 1217–1229. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
37. Dietz V. Proprioception and locomotor disorders. *Nat Rev Neurosci* 2002; 10: 781–790. | [Article](#) | [PubMed](#) | [ISI](#) |
38. Dietz V, Muller R, Colombo G. Locomotor activity in spinal man: significance of afferent input. *Brain* 2002; 125: 2626–2634. | [Article](#) | [PubMed](#) | [ISI](#) |
39. Wolpaw JR. *Spinal Cord Plasticity: Alterations in Reflex Function*. Kluwer Academic Publishers, 1997.
40. Holler M, Hopf HC. Posttraumatic synkinesis of the diaphragm and muscles of the brachial plexus. *Neurology* 1968; 19: 141–153. | [ChemPort](#) |
41. Robinson PK. Associate movements between limb and respiratory muscles as a sequel to brain injury. *Johns Hopkins Hosp* 1951; 89: 21–29. | [PubMed](#) | [ChemPort](#) |
42. Kew JJ *et al.* Abnormal access of axial vibratory input to deafferented somatosensory amputees. *J Neurophysiol* 1997; 77: 2753–2764. | [PubMed](#) | [ISI](#) | [ChemPort](#) |
43. Moore CI *et al.* Referred phantom sensation and cortical reorganisation after spinal cord injury. *J Neurosci* 2000; 20: 14703–14708. | [Article](#) | [PubMed](#) | [ChemPort](#) |
44. Banati RB. Brain plasticity and microglia: is transsynaptic glial activation in the thalamus cortical plasticity and central sensitisation? *J Physiol Paris* 2002; 96: 289–299. | [Article](#) | [PubMed](#) |
45. Jain N, Florence SL, Kaas JH. Reorganisation of somatosensory cortex after nerve and spinal cord injury. *Neurosci Lett* 1998; 243: 143–149. | [PubMed](#) |
46. Moore KA, Kohno T, Karchewski LA, Scholz J, Baba H, Woolf CJ. Partial peripheral nerve transection facilitates GABAergic inhibition in the superficial dorsal horn of the spinal cord. *J Neurosci* 2002; 22: 6731. | [PubMed](#) | [ISI](#) | [ChemPort](#) |
47. Turton AJ, Butler SR. Referred sensations following stroke. *Neurocase* 2001; 7: 397–405.
48. Borsook D *et al.* Acute plasticity in the human somatosensory cortex following amputation. *J Neurosci* 2001; 21: 1017. | [PubMed](#) | [ChemPort](#) |
49. Kaas JH, Merzenich MM, Killackey HP. The reorganisation of somatosensory cortex following peripheral nerve injury in adult and developing mammals. *Ann Rev Neurosci* 1983; 6: 325–356. | [Article](#) | [PubMed](#) |
50. Vroemen M, Aigner L, Winkler J, Weidner N. Adult neural progenitor cell grafts survive and integrate along axonal pathways. *Eur J Neurosci* 2003; 18: 743–751. | [Article](#) | [PubMed](#) |
51. Hill CE *et al.* Acute transplantation of glial-restricted precursor cells into spinal cord contusion lesion. *J Neurosci* 2002; 22: 1017. | [Article](#) | [PubMed](#) | [ChemPort](#) |

- differentiation, and effects on lesion environment and axonal regeneration. *Exp Neurol* 2003; 181: 310. | [Article](#) | [PubMed](#) | [ChemPort](#) |
52. Hofstetter CP *et al.* Allodynia limits the usefulness of intraspinal neural stem cell grafts; d outcome. *Nat Neurosci* 2005; 8: 346–353. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
53. Akiyama Y, Radtke C, Kocsis JD. Remyelination of the rat spinal cord by transplantation c cells. *J Neurosci* 2002; 22: 6623–6630. | [PubMed](#) | [ChemPort](#) |
54. Hofstetter CP *et al.* Marrow stromal cells form guiding strands in the injured spinal cord a *Acad Sci USA* 2002; 99: 2199–2204. | [Article](#) | [PubMed](#) | [ChemPort](#) |
55. Ankeny DP, McTigue DM, Jakeman LB. Bone marrow transplants provide tissue protectio axons after contusive spinal cord injury in rats. *Exp Neurol* 2004; 190: 17–31. | [Article](#) | [PubMed](#) |
56. Lu P, Jones LL, Tuszynski MH. BDNF-expressing marrow stromal cells support extensive injury. BDNF-expressing marrow stromal cells support extensive axonal growth at sites of *Neurosci* 2005; 191: 344–360. | [Article](#) | [PubMed](#) | [ChemPort](#) |
57. Kamada T *et al.* Transplantation of bone marrow stromal cell-derived Schwann cells prom functional recovery after complete transection of adult rat spinal cord. *J Neuropathol Exp Neurol* 2005; 150: 45. | [PubMed](#) |
58. Neuhuber B, Timothy Himes B, Shumsky JS, Gallo G, Fischer I. Axon growth and recovery bone marrow stromal cells in the injured spinal cord exhibit donor variations. *Brain Res* 2005; 1050: 85. | [Article](#) | [PubMed](#) | [ChemPort](#) |
59. Barnett SC. Olfactory ensheathing cells: unique glial cell types? *Neurotrauma* 2004; 21: 37–47. | [Article](#) | [PubMed](#) | [ChemPort](#) |
60. Raisman G. Olfactory ensheathing cells and repair of brain and spinal cord injuries. *Cloning Stem Cells* 2005; 7: 368. | [Article](#) | [PubMed](#) | [ChemPort](#) |
61. Fouad K, Schnell L, Bunge MB, Schwab ME, Liebscher T, Pearse DD. Combining Schwann ensheathing glia grafts with chondroitinase promotes locomotor recovery after complete t *Neurosci* 2005; 25: 1169–1178. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
62. Blits B *et al.* Rescue and sprouting of motoneurons following ventral root avulsion and rei intraspinal adeno-associated viral vector-mediated expression of glial cell line-derived ne neurotrophic factor. *Exp Neurol* 2004; 189: 303–316. | [Article](#) | [PubMed](#) | [ChemPort](#) |
63. Ruitenberg MJ *et al.* Adeno-associated viral vector-mediated gene transfer of brain-derive atrophy of rubrospinal neurons following both acute and chronic spinal cord injury. *Neurosci Lett* 2005; 380: 406. | [Article](#) | [PubMed](#) | [ChemPort](#) |
64. Ruitenberg MJ, Hendriks WJT, Boer GJ, Verhaagen J. Gene therapy for spinal cord injury *Gene Therapy in the Central Nervous System*. Elsevier Inc.: Amsterdam 2006. pp 273–28
65. Gaunt RA, Prochazka A. Control of urinary bladder function with devices: successes and fa *Neurosci Biobehav Rev* 2004; 28: 163–194. | [PubMed](#) |
66. Prochazka A, Mushahwar VK, McCreery DB. Neural prostheses. *J Physiol* 2001; 533: 99–109. | [Article](#) | [PubMed](#) | [ChemPort](#) |
67. Balsubramanian AV, Wood S, Middleton FRI, Fowler C, Craggs MD. Urinary guarding refl *Br J Urol Int* 2004; 9: 4.
68. Balsubramanian AV, Craggs MD, Fowler CJ. A neurophysiological and urodynamic assess sensory–motor pathways and their influence on aberrant lumbar–sacral reflexes in incom *Res Rev* 2004; 9: 111–115.
69. Rodi Z, Vodusek DB. The sacral reflex studies:single versus double pulse electrical stimula *Neurosci Lett* 2004; 368: 496–497.
70. Curt A, Dietz V. Electrophysiological recordings in patients with spinal cord injury: signifi *Cord* 1999; 37: 157–165. | [Article](#) | [PubMed](#) | [ISI](#) | [ChemPort](#) |
71. Marshall LF *et al.* Deterioration following spinal cord injury. A multicenter study. *J Neuro* 2005; 127: 404. | [PubMed](#) | [ISI](#) | [ChemPort](#) |
72. Granovsky Y, Matre D, Sokolik A, Lorenz J, Casey KL. Thermoreceptive innervation of hu contact heat evoked potential analysis. *Pain* 2005; 115: 238–247. | [PubMed](#) |

73. Steeves J, Fawcett J, Tuszynski M. Report of international clinical trials workshop on spin 2004, Vancouver, Canada. Spinal Cord 2004; 42: 591–597. | [Article](#) | [PubMed](#) | [Chem](#)
74. Christopher Reeve Foundation. Can non-human primate experiments expedite translation interventions following spinal cord injury to humans? in revision.
75. Edgerton VR, Roy RR, Hodgson JA, Prober RJ, de Guzman CP, de Leon R. Potential of ad cord to execute and acquire improved locomotion in the absence of supraspinal input. J N S128. | [PubMed](#) |
76. Dietz V, Harkema SJ. Locomotor activity in spinal cord-injured persons. J Appl Physiol 20 1960. | [Article](#) | [PubMed](#) | [ChemPort](#) |

MORE ARTICLES LIKE THIS

These links to content published by NPG are automatically generated

NEWS AND VIEWS

A question of making it work

Nature News and Views (13 Jan 1994)

Overcoming inhibition

Nature News and Views (30 Nov 1995)

See all 11 matches for News And Views

RESEARCH

The injured spinal cord spontaneously forms a new intraspinal circuit in adult rats

Nature Neuroscience Article (01 Mar 2004)

See all 46 matches for Research

Top

Spinal Cord

ISSN: 1362-4393

EISSN: 1476-5624

- [About NPG](#)
- [Contact NPG](#)
- [Nature jobs.com](#)
- [Privacy policy](#)
- [Legal notice](#)
- [Accessibility statement](#)
- [RSS web feeds](#)
- [Help](#)

© 2007 International Spinal Cord Society

partner of AGORA, HINARI, CrossRef and COUNTER

CareCure Forums > News Forums > Health & Science News > Poulsen, et al. (2002). Gene therapy for spinal cord injury and disease [In Process Citation]

PDA

View Full Version : Poulsen, et al. (2002). Gene therapy for spinal cord injury and disease [In Process Citation]

Max

04-20-2002, 05:14 PM

Gene therapy for spinal cord injury and disease [In Process Citation]

Selected

J Spinal Cord Med 2002 Spring;25(1):2-9 (ISSN: 1079-0268)

Poulsen DJ; Harrop JS; During MJ

University of Montana, Department of Pharmaceutical Sciences, Missoula 59812-1552, USA.

Poulsen@selway.umt.edu.

An incomplete understanding of the pathological processes involved in neurodegeneration and dysfunction of spinal cord injuries and diseases makes these disorders difficult to treat. Repair of damaged or genetically impaired spinal cord also has been limited by the complexity, cellular heterogeneity, and relative inaccessibility of the tissue. Thus, therapeutic options for the treatment of either chronic spinal cord diseases such as amyotrophic lateral sclerosis or acute spinal cord injuries have been rather limited. Potential new therapeutic targets are being identified as our understanding of the molecular pathology involved in neural injury and regeneration increases. Recent advances in gene transfer techniques have made gene therapy a more realistic and viable strategy for the treatment of a broad range of spinal cord disorders. This review summarizes the current state of knowledge regarding the limitations and recent advances in gene therapy and potential application of this technology toward spinal cord injury and disease.

[This message was edited by Wise Young on May 04, 2002 at 12:09 PM.]

Wise Young

04-24-2002, 03:36 PM

Max, if you post articles or abstracts... can you enter a brief description and why you want to post it in the header, and the actual article with the URL address to the body of the message? Thanks. Wise.

perry

04-24-2002, 04:09 PM

i have just heard the same thing from a person at the reeve's foundation. the primates studies will lead the way. gene therapy has been around for over 25 years, and now beginning to show success in animals. max,dr.wise how can we find out more.....

perry

Wise Young

04-25-2002, 10:02 AM

Perry, in spinal cord injury, there are two issues. The first is identification of beneficial (and deleterious) gene expression that can and should be manipulated in order to improve recovery, regeneration, remyelination. There are currently many laboratories systematically studying animal spinal cord injury models to identify regeneration-associated genes (RAGs), pain-associated genes (PAGs), neuroprotection-associated genes (NAGs), myelination-associated genes (MAGs), etc. Once identified, the expression of these genes can then be boosted or blocked in the spinal cord.

The second issue is the mechanism of changing gene expression. Gene therapy today is being carried out in several ways:

- Transgenic - the genes of an egg or sperm are modified and the subsequent organism then has a knockout (deleted), knockin (inserted), or dominant negative (an interfering gene) added. This is currently not an option for adult.
- In vivo transfection - the gene is inserted into certain cells by virus, liposomes, or other vectors. The viral

method is the most efficient and popular at the present but got into trouble recently, particularly adenovirus (the common cold virus), because it initiated fatal inflammation in one patient (Jesse Gelsinger). Many companies have touted other non-viral means of inserting genes that are generally less efficient but presumably safer.

- Ex vivo transfection and implantation of transfected cell - specifically cells can be removed from the body, transfected so that they express certain gene products, and then implanted back into the body. Actually, the first use of this technology in the CNS was for spinal cord injury (Tuszynski, et al.)

An alternative and growing approach to manipulating gene expression is with drugs. A large number of drugs and factors are known to turn off and on certain genes. For example, the tetracycline antibiotics are known to turn on certain genes. Likewise, there are many so-called nuclear factors that go into a cell and turn on genes, i.e. NF-kappa B is the factor that turns on inflammatory genes.

There is really no magic about gene expression. We just have much more powerful tools that allow us to measure and manipulate gene expression. So, once we know which genes we want to turn on and which to turn off or modulate, it can and will be done. Unfortunately, as a recent report in the Wall Street Journal suggests, the human genome project has not yielded a huge number of treatments for the pharmaceutical industry.

Despite massive investments, the number of drugs that have been approved by the FDA has fallen in the past three years compared to previous years. The reason is that people were expecting knowledge of the human gene to produce new drugs. They had not realized that the knowledge has produced the possibility of many drugs but the same amount of work and information must be gathered about each candidate drug before it can be successfully taken to clinical trial. Thus, the investment did not reduce the expense or time in developing the drugs. It just increased the number of potential drug candidates.

One of the most interesting outcomes of the human genome project is that it has shown us how similar humans are to other animals. I suspect that primate experiments will not be essential for moving all therapies into clinical trials. Scientists are working very hard to develop surrogate measures, using human cell cultures (stem cells, etc.) that allow testing of therapies without doing as many large animal experiments. Of course, the FDA continues to require large animal safety (toxicity) studies before clinical trial but every effort is being made to reduce the number of animals required.

Wise.

vBulletin v3.5.3, Copyright ©2000-2007, Jelsoft Enterprises Ltd.



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[\[Sign In\]](#) [\[Regis\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for *ex vivo spinal cord injury* [Go](#) [Clear](#)

Limits Preview/Index History Clipboard Details

Note: Performing your original search, *ex vivo spinal cord injury*, in PubMed will retrieve 38 citations.

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 1

☐ 1: Clin Neurosci. 1995-1996;3(5):268-74.

[Links](#)

Ex vivo gene therapy for Alzheimer's disease and spinal cord injury.

Blesch A, Tuszynski M.

Department of Neurosciences-0608, University of California-San Diego, La Jolla 92093-0608, USA.

Gene transfer is a potential means of treating chronic neurologic disorders and injury related neural degeneration. One approach for transferring genes to the CNS is to genetically modify cells in vitro and then transplant the cells to the CNS. For example, fibroblasts can be infected with a replication-defective retrovirus expressing a transgene, and can then be transplanted into the brain or spinal cord, thereby providing neurotrophic factors and substrates for axonal growth and elongation. In this review we discuss the grafting of neurotrophic factor secreting autologous fibroblasts in the rat and primate CNS. NGF secreting grafts have been shown to prevent degeneration of cholinergic neurons in the basal forebrain of primates and to induce sprouting of sensory, motor, and noradrenergic neurites after spinal cord injury. These results suggest the potential usefulness of ex vivo gene transfer for the treatment of Alzheimer's disease and spinal cord injury.

PMID: 8914793 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort by Send to

Related Links

Grafts of fibroblasts genetically modified to secrete NGF, BDNF, NT-3, or basic FGF elicit differential responses in the adult primate CNS. [Cell Transplant. 1996]

Gene therapy in the adult primate brain: intraparenchymal grafts of cells genetically modified to produce nerve growth factor prevent cholinergic neuronal degeneration. [J Neurosci. 1996]

Functional characterization of NGF-secreting cell grafts to the acutely injured spinal cord. [Cell Transplant. 1997]

Maintaining the neuronal phenotype after injury in the adult CNS. Neurotrophic factors, axonal growth substrates, and gene therapy. [J Neurosci. 1995]

Nerve growth factor delivery by gene transfer induces differential outgrowth of sensory, motor, and noradrenergic neurites after adult spinal cord injury. [Exp Neurol. 1996]

[See all Related Articles...](#)

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jan 29 2007 05:15:30

[nature.com homepage](#)

- [Jump to main content](#)
- [Jump to navigation](#)

[Login](#)



- [My account](#)
- [E-alert sign up](#)

- [Subscribe](#)
- [Register](#)

- [Publications A-Z index](#)
- [Browse by subject](#)

Search [Advanced search](#)



Reviews

Nature Reviews Neuroscience 7, 628-643 (August 2006) | doi:10.1038/nrn1955

There is a [Corrigendum \(1 November 2006\)](#) associated with this article.

Focus on: [Nerve regeneration](#)

Therapeutic interventions after spinal cord injury

Sandrine Thuret^{1,4}, Lawrence D. F. Moon^{2,4} and Fred H. Gage³ [About the authors](#)

Abstract

Spinal cord injury (SCI) can lead to paraplegia or quadriplegia. Although there are no fully restorative rehabilitative, cellular and molecular therapies have been tested in animal models. Many of these clinical trials. Here, we review these potential therapies, with an emphasis on the need for reproducible efficacy. Individual therapies are unlikely to provide a panacea. Rather, we predict that combining

improvements in outcome after SCI. Basic scientific research should provide a rational basis for clinical therapies to different types of SCI.

- View **At a Glance**

Worldwide, an estimated 2.5 million people live with spinal cord injury (SCI), with more than 1: year (see **International Campaign for Cures of Spinal Cord Injury Paralysis** in Online restorative therapies for SCI as yet and so prevention (for example, effective seat belts, weapons the best medicine (see **Foundation for Spinal Cord Injury Prevention, Care and Cure** in significant impact on quality of life, life expectancy and economic burden, with considerable cost loss of income. In one study, quadriplegics ranked recovery of arm and hand function as a prior recovery of sexual function as most important (when measured against recovery of bladder/bowel **autonomic dysreflexia**, improving walking movements and trunk stability, regaining normal pain)¹. Therapies addressing these and other important priorities (such as recovery of cardiovascular, skeletomuscular properties, and reducing spasticity) have been reviewed elsewhere^{2, 3, 4, 5, 6}. Limb function, which is the focus of most ongoing animal studies and clinical trials for treatment

To identify therapies that are unambiguously safe and effective, the scientific and clinical SCI community must ensure that preclinical studies be reproduced by independent laboratories, and that clinical trials have include an a priori unambiguous definition of primary outcome measures and any intended stratification methods that are sensitive enough to detect potentially small increments in function^{7, 8, 9}. Several studies have been evaluated independently under contractual arrangements between the National Institute of Neurological Disorders and Stroke (NINDS) and several Facilities of Research Excellence for SCI (FORE-SCI; see NINDS Facilities of Research Excellence for SCI in Online links box), including the Miami Project to Cure Paralysis and the Reeve–Irwin Institute for Neurological Science. For information on clinical trials, readers are directed to governmental and international consensus documents (see **Translating promising strategies for spinal cord injury therapy** in Online links box)¹⁰.


Here, we stress various cellular and molecular strategies that are supported by more than one preclinical study and that result in functional improvements after SCI; many of these strategies have reached, or are being tested, in clinical trials. Some of the potential therapies described below might produce only small improvements, and a combination of therapies may be needed to improve everyday quality of life. Speciality journals and general audience media need to accurately report the safety and efficacy of potential therapies to avoid raising and then dashing the hopes of those affected by SCI, those carrying out research, or the general public.

Endogenous response to SCI

The normal architecture of the human spinal cord (**Fig. 1**) can be radically disrupted by injury. The outcome^{13, 14} and can result from contusion, compression, penetration or maceration of the spinal cord, including neurons, oligodendrocytes, astrocytes and precursor cells¹⁵ (**Fig. 2**), and any reorganization of the spinal cord, additional structure and function are lost through active secondary processes (including loss of oligodendrocytes and loss of myelin¹⁶). Demyelinated axons are observed up to a decade after injury, and some of which these axons survive unmyelinated or become remyelinated by central or peripheral myelin sheaths¹⁷. Resident and invading inflammatory cells (including neutrophils, microglia, macrophages) play a wide range of destructive and reparative roles¹⁹. SCI culminates in glial scarring, a multifactorial process involving astrocytes, glial progenitors, microglia and macrophages^{20, 21}, fibroblasts and Schwann cells¹⁷.

perpendicular to the neuraxis and appears impenetrable. The scar also contains secreted and trapped axon growth^{23, 24}. Progressive expansion of the injury across more than one segment (syringomyelia) over months or years, sometimes proving fatal.


Figure 1 | Intact spinal cord.



a | Schematic showing a sagittal view through the human CNS. b | Transverse section through the spinal cord showing the relationship between axonal tracts and grey matter. c | Cortical, brainstem and spinal axons project through the white matter of grey matter, which in turn send axons through the PNS to target organs, including muscle. Primary sensory axons project through the PNS to second order sensory neurons in the CNS grey matter, which, in turn, send axons through the dorsal columns to supraspinal regions. Oligodendrocytes myelinate ascending and descending axons.

- [High resolution image and legend \(38 KB\)](#)
- [Figures and tables index](#)
- [Download Power Point slide \(248 KB\)](#)

Figure 2 | Spinal cord after injury.




Schematic showing a sagittal view through a region of cervical spinal cord injury (SCI), depicting different types of injury. Many cells die immediately, as well as progressively, after SCI. Cysts usually form after penetrating injury, cells from the PNS often invade the injury site to form a connective tissue scar, and progenitor cells and microglia. Many ascending and descending axons are interrupted and fail to regenerate. Some axons form new circuits with motor neurons via interneurons. At the site of cyst formation, cysts that are formed from ependymal cells. Disconnected myelinated axon segments are phagocytosed and, if spontaneous remyelination occurs, largely by PNS Schwann cells, whereas denervated (non-spa

- **High resolution image and legend (39 KB)**
- **Figures and tables index**
- **Download Power Point slide (247 KB)**

In contrast to these destructive events, commonly observed pathological features do indicate some degree of recovery (Fig. 2). Whereas there is little or no neurogenesis in the injured spinal cord, proliferation in the ventral horn generates new precursor cells that exclusively differentiate into glial cells^{15, 26, 27, 28}. Lignified lesions might even be spanned by trabeculae containing axon sprouts^{25, 29}. Sprouting is largely regulated by growth-inhibitory molecular factors^{24, 30}, and few axons regenerate over long distances back to their original targets. However, neuroplasticity occurs that could contribute to limited compensatory reorganization. At the cortical level, circuits can bypass the lesion, including sprouting of injured corticospinal axons onto spared, lower motor neurons that increase connectivity with lumbar motor neurons^{33, 34}. Cortical sensorimotor areas can functionally compensate for the subcortical level, the rubrospinal system can reorganize and compensate for much of the function lost to injury³¹.

Therefore, although there is some spontaneous repair after CNS injury, it is incomplete. Further combination of effective and safe therapeutic interventions (**Fig. 3**).

Figure 3 | Injured spinal cord after combination treatments.



Schematic showing a sagittal view through injured cervical spinal cord after a hypothetical combi are filled by vascularized grafts and trabeculae are spared. Grafts provide remyelinating cells, ar regions and in intact spinal cord are neutralized using antibodies, peptides or enzymes. Grafted relay circuits or the regeneration of injured axons back to their original targets. Furthermore, re synapses to be stabilized and reverses muscle atrophy.

- **High resolution image and legend (39 KB)**
- **Figures and tables index**
- **Download Power Point slide (246 KB)**

Cellular therapeutic interventions

Cellular transplantation after SCI has several aims: to bridge any cysts or cavities; to replace dead neurons or myelinating cells); and to create a favourable environment for axon regeneration.

Transplantation of peripheral nerve. After SCI in adult rats, autologous transplants (support ingrowth of various axonal types but not supraspinal axons³⁵. Peripheral nerve grafts v therapies (including anti-inflammatory drugs, vertebral wiring, fibrin glue and acidic fibroblast with regeneration of supraspinal axons into, through and beyond grafts^{36, 37, 38, 39}.

A similar strategy has been tested in non-human primates after lateral spinal hemisection⁴⁰. No axons were detected but some spinal axons were found to have regenerated 4 months after injury. This approach is being tested in chronic, incomplete human SCI, with one peer-reviewed report of limited functional recovery in 10 patients who were investigated⁴¹. Anecdotally, this strategy has not proved successful in people with SCI. Much work remains to be done to determine whether therapies that involve peripheral nerve grafts can effectively improve outcome after human SCI.


Transplantation of Schwann cells. Schwann cells from peripheral nerves have been transplanted into injured nerves, either being injected as suspensions after contusion injury⁴² or implanted into channels containing either hemisection⁴³ or complete transection⁴⁴. After transection and implantation of Schwann cells, Schwann cell bodies near the grafts extend into these bridge grafts, become myelinated⁴⁴ and are electrophysiologically active. Motor axons do not leave grafts distally to reinnervate the host. After contusion and implantation of Schwann cells, sensory and spinal axons extend into grafts, and many are remyelinated⁴². Recovery of hind

some⁴², but not all⁴⁶, studies. Consequently, combination therapies have been evaluated. After regeneration of CNS axons beyond bridges has been reported in response to transplantation of (delivery of neurotrophins^{47, 48}, a steroid (methylprednisolone sodium succinate)⁴⁹ or olfactory

Human Schwann cells have also been transplanted into the transected spinal cord of rats with a rats, brainstem axons regenerated into grafts and spinal axons regenerated distal to grafts. Function reported, although weight-supported stepping was observed in only one rat⁵¹. Finding the most combination therapy involving Schwann cells remains crucial. One important step towards human safety and efficacy of transplanting autologous Schwann cells into non-human primates after SCI has been no peer-reviewed reports of clinical trials involving the transplantation of Schwann cells at

Transplantation of olfactory nervous system cells. Cells from the embryonic and adult have been transplanted after SCI. Indeed, porcine, primate and human cells are now being tested in models of SCI and demyelination^{53, 54, 55, 56, 57}. Functional recovery and/or CNS axon regeneration of olfactory nervous system-derived cells are transplanted immediately or up to 2 months after SCI^{64, 65}. After lateral cervical hemisection in adult rats, injection of cells from the olfactory bulb led to improved function and enhanced performance on a climbing task⁵⁸. These transplants might also prevent SCI and may enhance myelination after SCI⁶⁸, although whether OEG directly myelinate axons after SCI is unclear. Transplants of cells from the olfactory nervous system do not, however, promote CNS axon regeneration under all circumstances^{42, 67, 70, 71}. FORE-SCI re-assessment of delayed transplantation of olfactory bulb cells after transection of adult rat spinal cord failed to find any improvement in hindlimb function, although some improvement was found in caudal spinal cord tissue⁷².

Figure 4 | The olfactory nervous system.



Schematic of a sagittal section through the human head, showing the olfactory nervous system (the olfactory nervous system depicted in greater detail (inset). Stem cells at the base of the olfactory epithelium produce new neurons throughout life, which extend axons de novo to the olfactory bulb. These axons are wrapped in myelin as they pass through the lamina propria from olfactory mucosa and into the CNS via the cribriform plate. permission, from Ref. 275 © (1996) TM Higher Education Group.

- **High resolution image and legend (52 KB)**
- **Figures and tables index**
- **Download Power Point slide (260 KB)**

Transplants of cells derived from fetal olfactory bulbs or the adult mucosa have reportedly been used in over 400 humans in China, Portugal and Colombia^{8, 73, 74}. Many procedures do not meet international standards, and, because controls are not included and comprehensive follow-up studies have not been performed, the safety and efficacy of this intervention, although there are reports of improvements in motor and

independent case report describes rapid segmental improvement in a single patient classified as according to the American Spinal Injury Association (ASIA) Impairment Scale (**Box 1**) — who is quadriplegic⁷⁵. An additional seven patients have been independently assessed pre- and post-op (including meningitis), and no clinically useful improvements, being observed. The view was to recommend this procedure to patients⁷⁴.

Box 1 | The ASIA Impairment Scale

- **Full box**

Elsewhere, formal veterinary and human clinical trials using cells derived from the adult olfactory bulb are advancing. SCI in dogs often occurs as a result of road traffic accidents or intravertebral disc extrusion. To test potential therapeutics in a large, heterogeneous patient population with variable injury severity, olfactory bulb have been transplanted autologously into nine dogs after naturally occurring thoracolumbar events up to 2 years later⁷⁶. Some hindlimb function was recovered (including weight-supportive gait), and controls and blind testing are required in future trials.

In one **Phase I clinical trial** in humans, cells were collected from the adult human lamina propria and transplanted into the spinal cord of three patients with thoracic injuries that had occurred at least 6 months prior. Controls were included. No adverse consequences were reported in these patients after 1 year, and no significant improvements were reported: a 3-year follow-up study is planned. A large number of patients and sensitive testing will be required to rule out the possibility that functional tissue has been derived from the complete patients, particularly before applying this therapy to incomplete injuries.

It is necessary to establish whether there are conditions under which transplantation of cells from the olfactory bulb works reproducibly to promote plasticity, regeneration, remyelination, neuroprotection and/or resolution of issues to be resolved include the optimal source of cells (lamina propria versus olfactory bulb), and graft strategy (for example, injection of suspensions or transfer within cellular matrix). It will be necessary to determine whether enriching cultures for specific phenotypes of cells improves outcome^{58, 69, 78, 79}.

Transplantation of embryonic CNS tissue. After spinal cord transection in animal models, axons grow into the lesion site, a small number of host axons regenerate into the transplant but terminate at the border^{80, 81}. Small but significant functional recovery is observed in rats^{82, 83} and cats⁸⁴. This recovery is due to axon growth into, through and beyond grafts, and the authors suggest that it is instead caused by relay synapses, affording transmission of signals via transplanted neurons, which are innervated by proximal axons and turn to distal host neurons. Grafts might also provide growth factors or improve conduction in spinal cord transplants are combined with neurotrophin delivery after complete spinal cord transection, functional recovery is observed⁸⁷, with some supraspinal and propriospinal axons growing into the caudal spinal cord.

Intraspinal transplantation of fetal spinal cord has been tested in a clinical trial involving patients with syringomyelia. Complications were observed and cysts were obliterated in all the patients. These trials have not been followed by standard treatment for SCI or syringomyelia⁷, perhaps because of the difficulties associated with transplantation.

Transplantation of embryonic stem/progenitor cells. Multipotent progenitor cells can self-renew indefinitely and differentiate into any cell type. Three of the major challenges in repair after SCI are controlling the survival, integration and differentiation of transplanted cells


functional recovery by reconstituting damaged circuits, remyelinating axons, and increasing plaques. Many groups have studied the fate of stem cells^{91, 92} or progenitor cells^{93, 94, 95, 96, 97, 98, 99} derived from embryonic CNS or human umbilical cord blood and transplanted into the injured adult rodent. These studies have reported modest improvements in functional recovery^{91, 100, 101}.

The potential of human fetal stem cells in animal models of SCI is currently being investigated. In human fetuses, human fetuses have been transplanted into immunosuppressed mice¹⁰² and non-human primate cases, the transplanted cells survived and differentiated into cells with characteristics of oligodendrocytes associated with locomotor improvements^{102, 103}.

The most recent successful approach with embryonic CNS-derived stem/progenitor cells is to use pre-differentiated to a desired lineage before transplantation. Transplantation in rats of neuron-contusion injury improved bladder and motor function. The cells survived, filled the lesion site, and some characteristics of neurons and glia, resulting in sparing/sprouting of descending pathways. Transplantation of embryonic stem cell (ESC)-derived oligodendrocyte-restricted progenitor cells into the adult rat enhanced remyelination and promoted improvement of motor function. The cells survived, migrated, and differentiated into oligodendrocytes. By contrast, when cells were transplanted 10 months after injury, no remyelination or locomotor recovery^{105, 106}. This study is being considered for FORE-SCI replication. **Cure Paralysis** (see Online links box).

Transplantation of adult stem/progenitor cells. Adult stem cells are now being considered. In contrast to ESC transplantation, adult stem cell transplantation should reduce ethical concerns and should not be rejected. Various adult progenitor cells have been implanted in rodent models of SCI: olfactory system (see above) to bone marrow-derived stem cells, cultured spinal cord and brain cells¹⁰⁷ (Fig. 5).

Figure 5 | Potential sources of stem/progenitor cells for transplantation into the injured spinal cord



Stem/progenitor cells can be collected at three different stages of development: from the inner cell mass of the blastocyst; from the brain, spinal cord, olfactory system or umbilical cord of the fetus; and from the olfactory system, bone marrow or blood of the adult. Each of these cell populations can be propagated in culture to produce a molecule of interest, or be restricted to a particular cell fate before transplantation in the injured spinal cord. Some of these cells (those of fetal CNS origin and umbilical cord blood cells) could eventually be transplanted. Some of these cells have the potential to be used for autologous transplantation, including cells from umbilical cord blood cells (which can be frozen at birth for use in later life), haematopoietic stem cells. Also, endogenous stem/progenitor cells are present at the injury site and are actively dividing. Their fate and fate might provide an alternative to transplantation. This diagram is based on published data^{95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 108, 109, 110, 111, 113, 114, 115, 116, 117, 118, 119, 120}.

transplantation after SCI in animals.

- [**High resolution image and legend \(46 KB\)**](#)
- [**Figures and tables index**](#)
- [**Download Power Point slide \(254 KB\)**](#)

Box 2 | Ethics of spinal cord injury research and clinical trials

- [**Full box**](#)

Adult bone marrow contains several different stem cell populations, including haematopoietic stromal cells (BMSCs), also known as mesenchymal stem cells (**Box 3**). Transplantation of HSC after compression-induced SCI in mice^{108, 109} and transplantation of BMSCs significantly improved mice and rats^{109, 110, 111}. However, the potential mechanisms by which BMSCs act are currently unclear and axonal elongation-facilitating actions have been proposed¹¹⁰. Also, the functional outcomes are uncertain because many are primarily based on one evaluation protocol without other behavioural measures. A small-scale human trial was conducted in which autologous BMSCs were intravenously delivered. Improvements observed appeared to fall within an expected range of spontaneous recovery, and ASIA category B to D. Nevertheless, without controls or some indication of cell viability within the spinal cord, that a measure of procedural safety was demonstrated. To our knowledge, peer-reviewed results are limited to this study⁸.

Box 3 | Bone marrow cells

- [**Full box**](#)

Adult neural progenitor cells (NPCs), isolated from the dentate gyrus, the subventricular zone of the brain, are able to self-renew, and to be multipotent in vitro and after transplantation into the CNS^{112, 113}. After transplantation into the intact and injured murine spinal cord, differentiation into only astrocytes or oligodendrocytes was observed. Mouse brain-derived adult NPCs were transplanted into the injured spinal cord of adult rats. The growth factors to selectively increase the number of oligodendrocyte precursors after transplantation post-injury survived, migrated, integrated in the injured spinal cord tissue, generated mature oligodendrocytes, myelinated the injured axons, and promoted some functional recovery. However, NPCs transplanted 8 weeks after injury failed to exert similar effects¹¹⁷. Therefore, there is a need to find and neutralize the inhibitory factors that interfere with NPC survival after transplantation.

Adult neural stem cells also reside in the spinal cord¹¹⁸, and the ability to regulate their numbers and fate to provide an alternative to transplantation. To regulate their numbers and fate to promote recovery, which molecules are involved in governing neural stem cell proliferation, migration and differentiation. Endogenous stem cells would require no exogenous stem cell sources and would therefore circumvent immune rejection, as well as the ethical and moral considerations associated with their use.

Transplantation of engineered stem/progenitor cells. The injured adult spinal cord is characterized by poor survival, neuronal differentiation and maturation. Therefore, to enhance the capacity of stem cells to survive and differentiate, recently begun to engineer stem cells to have better survival, and desired differentiation and maturation. To attempt to increase the survival of transplanted rat ESCs, ESCs were genetically modified to overexpress survival factors.

protein. This led to tumour-like growth of cells, accompanied by increased morbidity and mortality. Transplanted in the compressed mouse spinal cord, engineered mouse ESCs expressing the cell adhesion molecule *NCAM* grew longer and migrated rostrally and caudally from the lesion. Corticospinal axons showed interdigitation and extended into and, in some cases, beyond the lesion site¹²⁰.

It is apparent that transplantation alone of stem/progenitor cells after SCI will not lead to optimum recovery. It will be necessary for optimum return of function. Advances in molecular biology (for example, viral-mediated manipulation of these cells to express molecules of interest^{121, 122}). These types of combination require further development and careful animal testing, individually and jointly, before any clinical trials.

Transplantation of activated macrophages. It has been suggested that the failure of the transplantation of activated macrophages is attributed to the nature of the macrophage response¹²³, which differs from that observed in the transplantation of macrophages. Improvement of hindlimb function has been reported after transection and transplantation of activated macrophages with PNS or skin tissue. Fibres were shown to extend through the lesion, and re-transection of the lesion restored previously recovered functions^{124, 125}. However, the degree of recovery was comparable to that achieved with other cell types and occurred only in a subgroup of rats¹²⁶.

By contrast, activation of intrinsic macrophages at the spinal contusion site with micro-injection of lipopolysaccharide had negative effects on hindlimb functional recovery and tissue survival¹²⁷. Depletion of macrophages with clodronate improved better hindlimb usage during overground locomotion, more extensive white matter sparing and

There is, therefore, evidence that macrophages have deleterious effects on functional recovery, and it would be advantageous to replicate these studies in independent laboratories. Moreover, no previous studies of transplantation of activated macrophages in non-human primates have been described.

Proneuron sponsored Phase I clinical trials on the transplantation of activated macrophages in the United States and Belgium (see **Proneuron** in Online links box). Blood-derived monocytes activated using bacterial lipopolysaccharide were transplanted into eight ASIA category A participants between 9 and 14 days after injury. No irreversible adverse effects were observed. All participants improved to ASIA category C, which was claimed to be well above the expected rate of improvement in a multicentre, randomized controlled, **Phase II clinical trial** for ASIA category A participants in the United States and Israel, but recruitment for this clinical trial has currently been suspended for financial reasons (see **Proneuron** in Online links box).

Molecular therapeutic interventions

Molecular therapies after SCI have several aims: to protect neurons from secondary cell death; to enhance conduction.

Neuroprotective therapies. Substantial effort has been devoted to limiting the evolution of the injury and development of neuroprotective measures for acute SCI (and, potentially, for accompanying surgery¹³⁰). Delivery of antibodies against a cell adhesion molecule present on neutrophils and monocytes reduced tissue damage after SCI in rats and improves motor function while reducing both autonomic dysreflexia and spasticity. Erythropoietin has been reported to improve outcome¹³⁰, although this finding might not be reproducible (see **Cure Paralysis** in Online links box). Several studies have also recently reported that intravenous administration of erythropoietin improves hindlimb function in mouse and rat models of SCI^{132, 133, 134}, and this common theme is being tested in clinical trials for SCI⁹.

Intravenous steroids (for example, methylprednisolone sodium succinate; MP) have been registered in many countries¹³⁵. There is considerable debate as to whether MP has been proved to be safe and effective^{137, 138, 139, 140}. Treatment is claimed, controversially, to be beneficial if an appropriate regimen of injury and whether more than 3 or 8 hours have elapsed since incurring the injury; however, treatment, incorrect dosing or treatment of penetrating SCI has been shown to be detrimental¹⁴². Randomized trials examined whether modest improvements have been shown using MP, GM-1 ganglioside (TRH), nimodipine and the NMDA (N-methyl-D-aspartate) antagonist gacyclidine⁸. In 7 trials, primary outcome measures were not significant and placebo controls were lacking. When observed, these were often based on post hoc stratification, and severe side effects were also reported. Trials of neuroprotective agents have shown that large multi-centre, double-blind studies for SCI placebo-controlled **Phase III clinical trials**, with primary outcomes clearly recorded a priori, highly effective and safe neuroprotective therapies for human SCI^{10, 11}.

Enhancing conduction. Electrophysiological studies of humans with chronic SCI indicate that demyelination and that only a proportion become remyelinated (although denuded axons might remyelinate (by host or transplanted glia) or enhance conduction could yet prove useful. A potent aminopyridine) that can improve axonal conduction has been tested in several double-blind, placebo-controlled trials with chronic SCI¹⁷. However, Acorda Therapeutics' Phase III clinical trials of an oral, sustained-release aminopyridine showed a trend for improvements only in spasticity (see **Acorda Therapeutics**).

Delivery of growth factors. Growth factors modulate neuronal survival, neurite outgrowth, neurotransmission. Exogenous administration of growth factors has been proposed as one potential approach. The effectiveness of this approach has been tested using, for example, brain-derived neurotrophic factor (BDNF)^{145, 146, 147}, basic fibroblast growth factor¹⁴⁸, glial cell-derived neurotrophic factor (GDNF)^{149, 150}, and neurotrophin 3 (NT3)^{141, 145, 146, 151, 152}, NT4 and NT5 (Ref. **153**). Growth factors can be delivered to the spinal cord by transient injection¹⁵⁴, continuous infusion^{143, 144} or insertion of an artificial carrier¹⁴². Ex vivo gene therapy involves grafting cells, usually fibroblasts, that have been transfected with growth factors^{145, 146, 151, 152, 153}. In vivo delivery of growth factors has also been achieved using adenovirus^{155, 156}, adeno-associated virus (AAV) and lentivirus¹⁵⁷ (see below).

After SCI, the exogenous delivery of NGF in rats can induce growth of coeruleospinal axons^{158, 159} and corticospinal axons¹⁶⁰. BDNF induces recovery of forelimb function after cervical lateral hemisection, rubrospinal, reticulospinal, vestibulospinal, raphespinal, and local sensory and motor axons¹⁶¹, and BDNF improves bladder and hindlimb function after a mid-thoracic contusion¹⁵², and GDNF promotes growth of dorsal column sensory axons after partial and complete spinal cord transections and induces recovery of function. Delivery of growth factors alone leads to only partial recovery, researchers are now combining these factors with other therapeutic approaches: OEG transplants and NT3 (Ref. **67**); marrow stroma-derived cells with serotonergic agonists and NT3 (Ref. **166**). In addition, delayed delivery of growth factors is necessary for acute delivery because axons of chronically injured neurons can lack appropriate growth factors.

Unfortunately, clinical trials using systemic delivery of growth factors for various disorders have shown no efficacy or unacceptable side effects, or both¹⁶⁸. Obviously, to avoid adverse effects, growth factors must be delivered in quantities to have an effect but their distribution must be restricted to the site at which they are delivered. In vivo NGF gene delivery in patients with Alzheimer's disease by implanting autologous fibroblasts expressing human NGF in the forebrain showed promising results, with no side effects attributable to the delivery.

of the rate of cognitive decline¹⁶⁹. However, to move forward with the clinical application of grc further work is required to show whether this promotes CNS axon regeneration and leads to fur human primates.

Delivery of cAMP or small GTPases. Cyclic AMP (cAMP) can induce axonal sprouting of c and of injured adult rat spinal sensory neurons in vivo when prophylactically applied^{171, 173, 174}. therapy needs to be effective when applied after SCI. In zebrafish, post-injury application of cAMP CNS axons and restored function¹⁷⁵. After injury, the CNS environment is more permissive for. Therefore, elevating cAMP levels after SCI has been tried in combination with other treatments. locomotion were observed^{176, 177} after delivery of Rolipram (which prevents the hydrolysis of c transplants¹⁷⁶, and after administration of the combination of Schwann cells, a cAMP analogue any human clinical tests can begin, therapeutic windows of delivery of cAMP analogues must be delivery established, ideally in contusion injury models in rodents or primates.

Other strategies targeting molecules that are intrinsic to neurons could be viable, with modulating approach. Many factors that limit axon regeneration (see below) signal to the neuronal cytoskeleton. Rho and Rac^{178, 179, 180}. Inhibition of Rho by a bacterial toxin, C3-ADP-ribosyltransferase, produced degree of functional recovery after dorsal hemisection injury in adult rats¹⁸¹, although these results study¹⁸². Side effects have also been reported^{182, 183} and, although potential explanations have efficacy of small GTPase modulation need to be further evaluated before their use for human SCI. Therapeutic has developed a cell-permeable variant of a Rho inhibitor known as Cethrin (BA-21 multi-centre Phase I/IIa trial that will include ASIA category A patients who are scheduled to receive days of thoracic SCI; Cethrin will be applied using fibrin¹⁸⁴ (see **BioAxone Therapeutic** in O

Rho kinase (ROCK) acts as a downstream effector of Rho¹⁸⁶. Inhibition of ROCK by a peptide-like molecule inhibitors stimulated or accelerated functional recovery, and had a neuroprotective effect in models when given locally or systemically immediately after injury either as a single dose or over ¹⁸⁸. However, it should be kept in mind that ROCK inhibitors have teratogenic potential¹⁸⁹ and functions of small GTPases might reduce the therapeutic specificity of the compounds that mod

Modulation of interactions with myelin inhibitors. Intact and injured CNS myelin contains molecules (including Nogo-A, myelin-associated glycoprotein, oligodendrocyte myelin glycoprotein B3)^{190, 191, 192, 193}. Various therapies have been developed to target and overcome these. delivery of anti-Nogo therapeutics, independent laboratories report CNS axon growth and recovery ^{195, 196, 197, 198, 199}, although not all^{200, 201}, rodent models of SCI, and report no nociceptive against Nogo-A have recently been shown to promote growth of corticospinal tract axons after contusion hemisection in four out of five marmoset monkeys tested¹⁹². Future experiments might show whether improve outcome in contusion or compression models of SCI. Phase I clinical trials using human delivery in progress for ASIA category A patients with thoracic SCI in association with Novartis (M. Schw

Therapies targeting molecules in receptor complexes for Nogo-A²⁰³ are also being tested. In some Nogo-A, Nogo receptor or NGF receptor leads to CNS axon growth and functional recovery²⁰⁴. factors in the negative studies need to be elucidated because these could be important future targets. delivery of **NgR(310)ecto-Fc** enhances corticospinal and raphespinal axon growth after dorsal contusion in rats and enhances electrophysiological and behavioural recovery^{209, 210}. Delayed, subcutaneous

promotes growth of corticospinal axons and serotonergic fibres and a degree of locomotor recovery after hemisection^{211, 212}: independent testing of NEP1-40 by one FORE-SCI centre is underway (see **Centre** in Online links box).

Extracellular matrix modifiers. Transient suppression of collagen synthesis promotes CN and, when combined with an analogue of cAMP, it has been reported to promote CNS axon regrowth after acute SCI²¹⁴ (but see Ref. ²¹⁵ for a contrasting result). Neuraxo has reported its intention combination therapy, which they have designated Cordaneurin, in human SCI (see **Neuraxo B** links box). However, it would be valuable to reproduce these results independently, and to carry out studies in primates.

In adult rats, degradation of growth-inhibitory chondroitin sulphate by delivery of the bacterial (ChABC) promotes regeneration of injured CNS axons and recovery of function after dorsal column spinal cord hemisection in adult rats, delivery of ChABC promotes regrowth of axons from spinal nerve grafts²¹⁸ and regrowth of CNS axons into the spinal cord beyond hemichannel bridges after complete transection and implantation of channels containing Schwann cells, delivery of ChABC promotes serotonergic axons beyond grafts²²⁰. Intrathecal delivery of ChABC also promotes recovery of locomotion following severe (although not moderate) thoracic contusion injury in adult rats²²¹. Tests for efficacy in human primate models of SCI remain to be reported. Seikagaku is testing ChABC in Phase II clinical trials (see **Seikagaku Corporation** in Online links box), which could aid translation to treatment.

Rehabilitative training

Improved locomotor function is often seen in mammals with incomplete and even complete SCI after rehabilitation²²². Locomotor training even enhances the ability of many spinally transected mammals to support body-weight support is provided^{31, 223, 224}. This improvement occurs because, after SCI, the spinal cord does not become silent but maintains active and functional neuronal properties, and can respond to the level of the injury. It can generate oscillating coordinated motor patterns and is capable of controlling locomotion. Increasing numbers of animal experiments combine rehabilitation/physical therapy with other approaches to promote regeneration and recovery of limb function^{228, 229, 230, 231}.

Many SCI clinical trials that are currently recruiting participants or are already in progress address locomotor training, including upper-extremity exercise, body-weight-supported treadmill training, robotic or manual locomotor training, and **functional electrical stimulation** (FES) (see **Clinical Trials.gov** in Online links box)². Studies have empirically shown which types of locomotor training and rehabilitation are optimal for recovery of function. Locomotor training enhances the ability of humans with neurologically complete SCI to walk on a treadmill when body-weight support is provided^{31, 223, 224}, although rehabilitation does not yet enable patients with complete SCI to walk unassisted overground²³³. FES of the dorsal surface of the spinal cord can induce step-like locomotion and corresponding electromyographic activity in the leg muscles in patients with complete SCI²²⁶. A recent randomised controlled trial has shown that many patients with recent, incomplete SCI achieve independent walking either using conventional devices or using body-weight-supported treadmill training^{234, 235}. Patients also benefit from treadmill or overground locomotor training: for example, improvements are seen in walking speed (although outside the testing environment, participants did not walk independently of their wheelchairs). An ASIA category C patient reported that a combination of treadmill training and spinal cord epidural FES increased the quantity of stepping during the training session and resulted in an immediate improvement in walking speed.

superior to that obtained with only treadmill training²³⁶. Therefore, the combination of central peripherally (locomotor training) induced stepping appears to be an effective method for restoration of normal supraspinal input and should be explored further. Improvements in health have also including improved cardiovascular performance and reductions in spasticity, bone loss and bladder

The mechanisms by which physical therapy or rehabilitation improve function after SCI need to be understood to allow for rational improvement in therapy. Experimentation is also vital to identify safe and effective exercise. Exercise can pose special risks to people with SCI, including autonomic dysreflexia, fracture or falls. People with SCI have atypical physiological responses to exercise (for example, abnormal heart rate response to sustain intense exercise². Inappropriate exercise could also be detrimental after SCI^{238, 239}. Heart rate is a potential confounding factor in clinical trials because it is difficult to control, although we should proceed without strong justification.

Despite the documented advantages of exercise and rehabilitation, a US survey of quadriplegics reported having no access to exercise, and a further ~45% reported having to exercise on their own without a physical therapist¹. Therefore, much remains to be done politically to ensure that therapies that are made available to individuals with SCI.

Technical aspects

Translating cellular therapies to the clinic. Because autologous transplants of cells or tissues require immunosuppression to escape immune rejection, they represent an attractive therapeutic option. A source for autologous grafts of peripheral nerves or Schwann cells because only a minor deficit in olfactory mucosa biopsy. The olfactory mucosa is more accessible than the olfactory bulb for autologous transplant. Autologous transplants using tissue from the olfactory bulb have been carried out in dogs⁷⁶. Expression of neurotrophins⁵⁴ or other mitogens²⁴¹ might be possible when the amount of tissue is limiting, and proliferation after transplantation needs to be prevented²⁴².

Cellular suspensions can be transplanted into the acute, post-injury milieu or into irregularly shaped cavities later in the injured spinal cord. Tissue grafts (for example, peripheral nerve grafts) are perhaps better suited for anatomically complete injuries or for external routing (for example, direct routes of administration might include delivery of cells into the cerebrospinal fluid by lumbar puncture towards the injury and exert a beneficial effect by reducing injury size^{243, 244}. Lumbar puncture has its minimal invasiveness, simplicity and low cost.

Cells could also be genetically modified to deliver therapeutic molecules^{145, 146, 151, 152, 153, 156} and include fibroblasts, ESCs, neural stem/progenitor cells, OEG and Schwann cells. However, in most cases transplanted cells die after transplantation and are replaced by host cells^{245, 246, 247}. Although this might still confer benefits, ensuring survival of the cells and controlling regulation of expression are important for transgenic delivery. Identifying transplanted cells requires the use of a marker that neither induces nor transfers to host cells²⁴⁶.

The protective or reparative potential of transplants of a given cell type can be established only by comparing alternative cell types (rather than merely injections of fluid). With regard to complete injuries, comparison regardless of the cell type transplanted¹²⁶; a goal for the future (currently elusive) will be to ensure that the transplant supports body weight²²⁶. Finally, it might be short-sighted to select a cell type for a clinical trial

cell types within a single experiment. If the race to clinical trial results in one cell type becoming been evaluated against other cell types, then other (potentially better) cells might not be easily t difficult to deny a clinical trial participant a therapy that has already been shown to be partially case when evaluating potential drug alternatives to MP¹³⁸.

Translating molecular therapies to the clinic. Techniques to deliver molecular therapies: intracerebroventricular, intrathecal and intraspinal injection, continuous infusion or insertion c molecule of interest. Viral vector-mediated transfer of molecules to the injured spinal cord is en strategy¹⁵⁷. In vivo gene therapy has been tested in models of SCI using viruses, including herp lentivirus and Moloney leukaemia virus²⁴⁸. Particularly interesting is the finding that AAV, wh retrogradely transported efficiently to motor neurons of the spinal cord²⁴⁹. It is an efficient too factor 1 and it extends life expectancy in a murine model of motor neuron disease²⁴⁹. AAV-med paraplegin also rescued peripheral axonopathy in a model of hereditary spastic paraplegia²⁵⁰. I injections could be a method for delivering a therapeutic molecule after SCI. However, impleme research to determine the best AAV serotypes to target motor neurons efficiently, and retrograd tested in the context of SCI.

An opportunity exists for tailoring therapies to different types of injury. For example, if regener desired, knowledge of the receptors expressed on the cell body and axon will inform whether thi particular neurotrophin, and where this factor might best be applied. Similarly, there might be l that neutralizes a given inhibitory receptor if this molecule is not expressed by the axons that ar a rational basis for intervening with a given therapy by meticulously investigating the mechanis

Preclinical testing. Many preclinical therapies have not been shown to be safe and efficaciou Independent replication is extremely desirable to determine the general applicability of a therap potential therapies should be tested in models that closely approximate the human injury subty injuries in dogs, as well as surgically induced injuries in non-human primates, can be used adva response to SCI, although studied surprisingly little, has been examined after contusion injury⁴¹ differences between rodent, cat, dog and primate nervous systems¹²⁶, many recommend that th primates for safety and efficacy^{10, 11}. Despite the paucity of safety and efficacy studies using no studies and trials in humans are currently in progress⁸. This trend is of particular concern giver including transplants of stem cells or cells from the olfactory nervous system, can induce pain-r growth of sensory and sympathetic axons when tested in rodent models of SCI^{72, 110, 254, 255}. : nociception, autonomic dysreflexia and spasticity should therefore take place in animal models patients to ensure that therapies neither induce adverse consequences nor interfere with the nat function that can occur. For example, when transplanting cells, care should be taken not to abla trabeculae or axons spared in circumferential white matter^{29, 256}.

There are also relatively few studies that report outcomes after intervening more than 1 month I ²⁵⁸, and, of these, many fail to detect improvements in axon growth or functional recovery. This repair is to be achieved in individuals with long-standing injuries. Additionally, relatively few st functional recovery go on to determine whether these changes remain stable beyond 2 or 3 mon

Clinical trials networks. Various databases of patients with SCI have been established to fo SCI and to enlist and document patients that might be suitable for particular clinical trials. Eurc trial networks have been established to be ready to implement interventions across multiple cer

standardized evaluation using clinical outcome measures, imaging and neurophysiological stimulation. Researchers and others, including the FORE-SCI groups, are developing additional tests of sensory function to allow more sensitive assessment of recovery of function after SCI⁷.

Conclusions

SCI is a devastating condition for which there is as yet no cure. Cellular, molecular and rehabilitative therapies have been developed and some are now in, or moving towards, clinical trials. Nevertheless, work remains to be done to determine if any of these therapies can safely improve outcome after human SCI. To distinguish therapies that are likely to be effective, the scientific and clinical SCI communities recommend that preclinical studies should be reproducible and that individual therapies are unlikely to emerge as a cure for SCI. Rather, we predict that tailored combinations of therapies will allow cumulative improvements in outcome after different types of SCI.

Links

FURTHER INFORMATION

- [Acorda Therapeutics](#)
- [American Spinal Injury Association](#)
- [BioAxone Therapeutic](#)
- [Christopher Reeve Foundation](#)
- [Clinical Trials.gov](#)
- [Foundation for Spinal Cord Injury Prevention, Care and Cure](#)
- [International Campaign for Cures of Spinal Cord Injury Paralysis](#)
- [Miami Project to Cure Paralysis \(FORE-SCI\)](#)
- [National Institute of Neurological Disorders and Stroke \(NINDS\) Facilities of Spinal Cord Injury](#)
- [Neuraxo Biopharmaceuticals](#)
- [NINDS workshop on translating promising strategies for spinal cord injury treatment](#)
- [Proneuron](#)
- [Reeve-Irvine Research Centre, University of California, Irvine \(FORE-SCI\)](#)
- [Seikagaku Corporation](#)

Acknowledgements

We thank the Christopher Reeve Foundation and the members of their International Research Consortium for their support. We also thank the Associate members of this Consortium while working in the laboratories of F.H.G. and M. B. Bunge. We thank L. Gage for editing the manuscript. Discussion of potential therapies in this review does not constitute an endorsement.

Competing interests statement

The authors declare **competing financial interests**.

References

1. Anderson, K. D. Targeting recovery: priorities of the spinal cord-injured population. *J. Neurosci.* 27, 11911–11912 (2007). **An important study that reveals the factors that patients with SCI consider to enhance their quality of life.**

- **Article**
- **PubMed**
- **ISI**
- 2. Nash, M. S. Exercise as a health-promoting activity following spinal cord injury. *J. Neurol* (2005).
- 3. Widerstrom-Noga, E. G. & Turk, D. C. Types and effectiveness of treatments used by people with spinal cord injuries: influence of pain and psychosocial characteristics. *Spinal Cord* 41, 60 (2003).
- **Article**
- **ChemPort**
- 4. Waxman, S. G. & Hains, B. C. Fire and phantoms after spinal cord injury: Na⁺ channels and pain. *J. Neurosci* 26, 207–215 (2006).
- **Article**
- **ChemPort**
- 5. Eaton, M. J. Cell and molecular approaches to the attenuation of pain after spinal cord injury. *J. Neurosci* 26, 1200–1208 (2006).
- **Article**
- 6. Kirshblum, S. Treatment alternatives for spinal cord injury related spasticity. *J. Spinal Cord Med* 29, 10–18 (2006).
- **PubMed**
- **ChemPort**
- 7. Kleitman, N. Keeping promises: translating basic research into new spinal cord injury therapies. *J. Neurosci* 24, 318–325 (2004).
- **PubMed**
- 8. Amador, M. J. & Guest, J. D. An appraisal of ongoing experimental procedures in human spinal cord injury. *Spinal Cord* 29, 70–86 (2005).
- 9. Blight, A. R. & Tuszynski, M. H. Clinical trials in spinal cord injury. *J. Neurotrauma* 23, 581–590 (2006).
- **Article**
- 10. Anderson, D. K. *et al.* Recommended guidelines for studies of human subjects with spinal cord injury. *Spinal Cord* 43, 458–465 (2005).
- **Article**
- **PubMed**
- **ISI**
- **ChemPort**
- 11. Steeves, J. , Fawcett, J. & Tuszynski, M. Report of international clinical trials workshop on spinal cord injury, 2004, Vancouver, Canada. *Spinal Cord* 42, 591–597 (2004).
- **Article**
- **PubMed**
- **ChemPort**
- 12. Hall, E. D. Drug development in spinal cord injury: what is the FDA looking for? *J. Rehabil. Med* 38, 10–18 (2006).

13. Bunge, R. P. , Puckett, W. R. , Becerra, J. L. , Marcillo, A. E. & Quencer, R. M. Observation a review and classification of 22 new cases with details from a case of chronic cord compres demyelination. Adv. Neurol. 59, 75–90 (1993).
 - o [PubMed](#)
 - o [ChemPort](#)
14. Kakulas, B. A. A review of the neuropathology of human spinal cord injury with emphasis Med. 22, 119–124 (1999).
 - o [PubMed](#)
 - o [ChemPort](#)
15. Horky, L. , Galimi, F. , Gage, F. H. & Horner, P. J. Fate of endogenous stem/progenitor cel Comp. Neurol. (in the press).
16. Crowe, M. J. , Bresnahan, J. C. , Shuman, S. L. , Masters, J. N. & Beattie, M. S. Apoptosis a spinal cord injury in rats and monkeys. Nature Med. 3, 73–76 (1997).
 - o [Article](#)
17. Guest, J. D. , Hiester, E. D. & Bunge, R. P. Demyelination and Schwann cell responses adj: following chronic human spinal cord injury. Exp. Neurol. 192, 384–393 (2005).
 - o [Article](#)
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
18. Totoiu, M. O. & Keirstead, H. S. Spinal cord injury is accompanied by chronic progressive 486, 373–383 (2005).
 - o [Article](#)
 - o [PubMed](#)
 - o [ISI](#)
19. Jones, T. B. , McDaniel, E. E. & Popovich, P. G. Inflammatory-mediated injury and repair cord. Curr. Pharm. Des. 11, 1223–1236 (2005).

Starts with an excellent primer explaining the body's innate and adaptive imr focus on the detrimental and potential reparative roles of neutrophils, macro cells.

 - o [Article](#)
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
20. Jones, L. L. , Yamaguchi, Y. , Stallcup, W. B. & Tuszynski, M. H. NG2 is a major chondroit after spinal cord injury and is expressed by macrophages and oligodendrocyte progenitors (2002).
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
21. Jones, L. L. , Margolis, R. U. & Tuszynski, M. H. The chondroitin sulfate proteoglycans ne

versican are differentially regulated following spinal cord injury. *Exp. Neurol.* 182, 399–411 (2002).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

22. Bruce, J. H. *et al.* Schwannosis: role of gliosis and proteoglycan in human spinal cord injury. *Neurosci. Lett.* 285, 141–144 (2000).

- [ChemPort](#)

23. Silver, J. & Miller, J. H. Regeneration beyond the glial scar. *Nature Rev. Neurosci.* 5, 146–157 (2004).

- [Article](#)

24. Fawcett, J. W. Overcoming inhibition in the damaged spinal cord. *J. Neurotrauma* 23, 371–384 (2006).

- [Article](#)

25. Beattie, M. S. *et al.* Endogenous repair after spinal cord contusion injuries in the rat. *Exp. Neurol.* 172, 115–127 (2001).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

26. Yang, H. *et al.* Endogenous neurogenesis replaces oligodendrocytes and astrocytes after spinal cord injury. *Neurosci.* 26, 2157–2166 (2006).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

27. Yamamoto, S., Yamamoto, N., Kitamura, T., Nakamura, K. & Nakafuku, M. Proliferation in response to injury in the adult rat spinal cord. *Exp. Neurol.* 172, 115–127 (2001).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

28. Azari, M. F., Profyris, C., Zang, D. W., Petratos, S. & Cheema, S. S. Induction of endogenous models of spinal cord injury and disease. *Eur. J. Neurol.* 12, 638–648 (2005).

- [Article](#)
- [ChemPort](#)

29. Hill, C. E., Beattie, M. S. & Bresnahan, J. C. Degeneration and sprouting of identified descending axons after contusive spinal cord injury in the rat. *Exp. Neurol.* 171, 153–169 (2001).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

30. Pettigrew, D. B., Shockley, K. P. & Crutcher, K. A. Disruption of spinal cord white matter : axonal growth *in vitro* in the absence of glial scarring. *BMC Neurosci.* 2, 8 (2001).

- [Article](#)

- [PubMed](#)
 - [ChemPort](#)
- 31. Raineteau, O. & Schwab, M. E. Plasticity of motor systems after incomplete spinal cord injury. *Neurosci. Lett.* 273 (2001).
- [Article](#)
- 32. Weidner, N. , Ner, A. , Salimi, N. & Tuszynski, M. H. Spontaneous corticospinal axonal sprouting after adult central nervous system injury. *Proc. Natl Acad. Sci. USA* 98, 3513–3518 (2001).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
- 33. Bareyre, F. M. *et al.* The injured spinal cord spontaneously forms a new intraspinal circuit. *Nat. Neurosci.* 7, 269–277 (2004).
Presents the important finding that significant spontaneous axonal sprouting occurs after spinal cord injury, which provides alternative routes through which corticospinal axons can reach the motor cortex.
- [Article](#)
- 34. Raineteau, O. , Fouad, K. , Bareyre, F. M. & Schwab, M. E. Reorganization of descending corticospinal tracts after spinal cord injury. *Eur. J. Neurosci.* 16, 1761–1771 (2002).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
- 35. Richardson, P. M. , McGuinness, U. M. & Aguayo, A. J. Axons from CNS grafts regenerate and form synapses in the adult rat spinal cord. *Neurosci. Lett.* 265 (1980).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 36. Cheng, H. , Cao, Y. & Olson, L. Spinal cord repair in adult paraplegic rats: partial restoration of locomotor function. *Neurosci. Lett.* 273, 510–513 (1996).
- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 37. Cheng, H. *et al.* Gait analysis of adult paraplegic rats after spinal cord repair. *Exp. Neurol.* 177, 1–11 (2002).
- [Article](#)
 - [ChemPort](#)
- 38. Lee, Y. S. , Hsiao, I. & Lin, V. W. Peripheral nerve grafts and aFGF restore partial hindlimb function after spinal cord injury. *Neurotrauma* 19, 1203–1216 (2002).
- [Article](#)
- 39. Lee, Y. S. , Lin, C. Y. , Robertson, R. T. , Hsiao, I. & Lin, V. W. Motor recovery and anatomical correlates in spinal cord-repaired adult rats. *J. Neuropathol. Exp. Neurol.* 63, 233–245 (2004).
- 40. Levi, A. D. *et al.* Peripheral nerve grafts promoting central nervous system regeneration after spinal cord injury. *Neurosci. Lett.* 368, 1–4 (2004).

primate. *J. Neurosurg.* 96, 197–205 (2002).

41. Cheng, H. , Liao, K. K. , Liao, S. F. , Chuang, T. Y. & Shih, Y. H. Spinal cord repair with acic treatment for a patient with chronic paraplegia. *Spine* 29, E284–E288 (2004).

- [Article](#)

42. Takami, T. *et al.* Schwann cell but not olfactory ensheathing glia transplants improve hind moderately contused adult rat thoracic spinal cord. *J. Neurosci.* 22, 6670–6681 (2002).

An important study that compares two competing cell types in a single experi

- [PubMed](#)

- [ISI](#)

- [ChemPort](#)

43. Bamber, N. I. *et al.* Neurotrophins BDNF and NT-3 promote axonal re-entry into the dista cell-seeded mini-channels. *Eur. J. Neurosci.* 13, 257–268 (2001).

- [Article](#)

- [PubMed](#)

- [ISI](#)

- [ChemPort](#)

44. Xu, X. M. , Guenard, V. , Kleitman, N. & Bunge, M. B. Axonal regeneration into Schwann c grafted into transected adult rat spinal cord. *J. Comp. Neurol.* 351, 145–160 (1995).

- [Article](#)

- [PubMed](#)

- [ISI](#)

- [ChemPort](#)

45. Pinzon, A. , Calancie, B. , Oudega, M. & Noga, B. Conduction of impulses by axons regener transected adult rat spinal cord. *J. Neurosci. Res.* 64, 533–541 (2001).

- [Article](#)

- [ChemPort](#)

46. Pearse, D. D. *et al.* Transplantation of Schwann cells and olfactory ensheathing glia after s with methylprednisolone and interleukin-10 enhance recovery? *J. Neurotrauma* 21, 1223–

- [Article](#)

- [PubMed](#)

47. Menei, P. , Montero-Menei, C. , Whittemore, S. R. , Bunge, R. P. & Bunge, M. B. Schwann human BDNF promote enhanced axonal regrowth across transected adult rat spinal cord. (1998).

- [Article](#)

- [PubMed](#)

- [ISI](#)

- [ChemPort](#)

48. Xu, X. M. , Guenard, V. , Kleitman, N. , Aebischer, P. & Bunge, M. B. A combination of BD axonal regeneration into Schwann cell grafts in adult rat thoracic spinal cord. *Exp. Neurol*

- [Article](#)

- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
49. Chen, A. , Xu, X. M. , Kleitman, N. & Bunge, M. B. Methylprednisolone administration in Schwann cell grafts in transected adult rat thoracic spinal cord. *Exp. Neurol.* 138, 261–276 (1996).
- [Article](#)
 - [ChemPort](#)
50. Ramon-Cueto, A. , Plant, G. W. , Avila, J. & Bunge, M. B. Long-distance axonal regeneration in the spinal cord is promoted by olfactory ensheathing glia transplants. *J. Neurosci.* 18, 3803–3815 (1998).
- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
51. Guest, J. D. *et al.* The ability of cultured human Schwann cells within PAN/PVC guidance channels to promote axonal regeneration in the transected nude rat spinal cord. *Can. J. Neurol. Sci.* 23, S1–S2 (1996).
52. Iwanami, A. *et al.* Establishment of graded spinal cord injury model in a nonhuman primate. *Neurosci. Res.* 80, 172–181 (2005).
- [Article](#)
 - [ChemPort](#)
53. Imaizumi, T. , Lankford, K. L. , Burton, W. V. , Fodor, W. L. & Kocsis, J. D. Xenotransplantation of olfactory ensheathing cells promotes axonal regeneration in rat spinal cord. *Nature Biotechnol.* 18, 105–111 (2000).
- [Article](#)
54. Bianco, J. I. , Perry, C. , Harkin, D. G. , Mackay-Sim, A. & Feron, F. Neurotrophin 3 promotes axonal regeneration in the rat spinal cord by recruiting olfactory ensheathing cells from human nose. *Glia* 45, 111–123 (2004).
- [Article](#)
55. Kato, T. , Honmou, O. , Uede, T. , Hashi, K. & Kocsis, J. D. Transplantation of human olfactory ensheathing cells promotes axonal regeneration and remyelination of demyelinated rat spinal cord. *Glia* 30, 209–218 (2000).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
56. Barnett, S. C. *et al.* Identification of a human olfactory ensheathing cell that can effect transected axonal regeneration in the demyelinated CNS axons. *Brain* 123, 1581–1588 (2000).
- [Article](#)
 - [PubMed](#)
57. Casas, C. E. *et al.* Non-human primate model of autologous olfactory ensheathing glia transplantation after electrolytic unilateral lesion of the medullary pyramid. *Abstr. Soc. Neurosci.* 184, 5 (2004).
58. Li, Y. , Decherchi, P. & Raisman, G. Transplantation of olfactory ensheathing cells into spinal cord promotes axonal regeneration and climbing. *J. Neurosci.* 23, 727–731 (2003).
- [PubMed](#)

- **ISI**
- **ChemPort**
- 59. Lu, J. , Feron, F. , Ho, S. M. , Mackay-Sim, A. & Waite, P. M. Transplantation of nasal olfactory ensheathing cells promotes recovery in paraplegic adult rats. *Brain Res.* 889, 344–357 (2001).
- **Article**
- **ChemPort**
- 60. Li, Y. , Field, P. M. & Raisman, G. Regeneration of adult rat corticospinal axons induced by olfactory ensheathing cells. *J. Neurosci.* 18, 10514–10524 (1998).
- **PubMed**
- **ISI**
- **ChemPort**
- 61. Li, Y. , Field, P. M. & Raisman, G. Repair of adult rat corticospinal tract by transplants of olfactory ensheathing cells. *Neuron* 27, 2000–2002 (1997).
- **Article**
- **PubMed**
- **ISI**
- **ChemPort**
- 62. Ramon-Cueto, A. , Cordero, M. I. , Santos-Benito, F. F. & Avila, J. Functional recovery of corticospinal axons after spinal cord transection by olfactory ensheathing glia. *Neuron* 25, 425–435 (2000).
- **Article**
- **PubMed**
- **ISI**
- **ChemPort**
- 63. Plant, G. W. , Christensen, C. L. , Oudega, M. & Bunge, M. B. Delayed transplantation of olfactory ensheathing cells promotes sparing/regeneration of supraspinal axons in the contused adult rat spinal cord. *J. Neurotrauma* 19, 149–161 (2002).
- **Article**
- **PubMed**
- **ISI**
- **ChemPort**
- 64. Lu, J. , Feron, F. , Mackay-Sim, A. & Waite, P. M. Olfactory ensheathing cells promote local axonal growth after transplantation into transected spinal cord. *Brain* 125, 14–21 (2002).
- **Article**
- **PubMed**
- **ISI**
- **ChemPort**
- 65. Keyvan-Fouladi, N. , Raisman, G. & Li, Y. Functional repair of the corticospinal tract by de novo differentiation of olfactory ensheathing cells in adult rats. *J. Neurosci.* 23, 9428–9434 (2003).
- **PubMed**
- **ChemPort**
- 66. Sasaki, M. , Hains, B. C. , Lankford, K. L. , Waxman, S. G. & Kocsis, J. D. Protection of corticospinal axons after spinal cord transection and engraftment of olfactory ensheathing cells. *Glia* 53, 352–359 (2006).
- **Article**
- 67. Ruitenberg, M. J. *et al.* NT-3 expression from engineered olfactory ensheathing glia promotes axonal growth in the injured spinal cord. *Neuron* 51, 101–114 (2006).

regeneration. *Brain* 128, 839–853 (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)

68. Sasaki, M. , Lankford, K. L. , Zemedkun, M. & Kocsis, J. D. Identified olfactory ensheathing transected dorsal funiculus bridge the lesion and form myelin. *J. Neurosci.* 24, 8485–849;

- [Article](#)
- [ChemPort](#)

69. Boyd, J. G. , Doucette, R. & Kawaja, M. D. Defining the role of olfactory ensheathing cells following damage to the spinal cord. *FASEB J.* 19, 694–703 (2005).

A thoughtful review that examines evidence for and against myelination by tr derived from the olfactory nervous system.

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

70. Riddell, J. S. , Enriquez-Denton, M. , Toft, A. , Fairless, R. & Barnett, S. C. Olfactory enshe influence on regeneration at the dorsal root entry zone following rhizotomy. *Glia* 47, 150–

- [Article](#)
- [PubMed](#)
- [ISI](#)

71. Ramer, L. M. , Richter, M. W. , Roskams, A. J. , Tetzlaff, W. & Ramer, M. S. Peripherally-d do not promote primary afferent regeneration following dorsal root injury. *Glia* 47, 189–2

- [Article](#)

72. Steward, O. *et al.* A re-assessment of the consequences of delayed transplantation of olfact complete spinal cord transection in rats. *Exp. Neurol.* 198, 483–499 (2006).

- [Article](#)

73. Huang, H. *et al.* Influence of patients' age on functional recovery after transplantation of c injured spinal cord injury. *Chin. Med. J. (Engl.)* 116, 1488–1491 (2003).

74. Dobkin, B. H. , Curt, A. & Guest, J. Cellular transplants in China: observational study from chronic spinal cord injury. *Neurorehabil. Neural Repair* 20, 5–13 (2006).

Independent evaluation of participants who had been transplanted with cells nervous system.

- [PubMed](#)

75. Guest, J. , Herrera, L. P. & Qian, T. Rapid recovery of segmental neurological function in a transplantation of fetal olfactory bulb-derived cells. *Spinal Cord* 44, 135–142 (2006).

- [Article](#)
- [ChemPort](#)

76. Jeffery, N. D. , Lakatos, A. & Franklin, R. J. Autologous olfactory glial cell transplantation occurring canine spinal cord injury. *J. Neurotrauma* 22, 1282–1293 (2005).

Describes safe veterinary intervention for SCI in dogs.

- [Article](#)
- [PubMed](#)
- [ISI](#)

77. Feron, F. *et al.* Autologous olfactory ensheathing cell transplantation in human spinal cord (2005).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

78. Lakatos, A. , Smith, P. M. , Barnett, S. C. & Franklin, R. J. Meningeal cells enhance limited olfactory ensheathing cells. *Brain* 126, 598–609 (2003).

- [Article](#)
- [PubMed](#)
- [ISI](#)

79. Murrell, W. *et al.* Multipotent stem cells from adult olfactory mucosa. *Dev. Dyn.* 233, 496–

- [Article](#)

80. Bregman, B. S. , McAtee, M. , Dai, H. N. & Kuhn, P. L. Neurotrophic factors increase axon: and transplantation in the adult rat. *Exp. Neurol.* 148, 475–494 (1997).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

81. Jakeman, L. B. & Reier, P. J. Axonal projections between fetal spinal cord transplants and neuroanatomical tracing study of local interactions. *J. Comp. Neurol.* 307, 311–334 (1991)

- [Article](#)
- [ChemPort](#)

82. Kunkel-Bagden, E. & Bregman, B. S. Spinal cord transplants enhance the recovery of locomotor function after spinal cord injury at birth. *Exp. Brain Res.* 81, 25–34 (1990).

- [ChemPort](#)

83. Bregman, B. S. *et al.* Recovery of function after spinal cord injury: mechanisms underlying locomotor function differ after spinal cord injury in newborn and adult rats. *Exp. Neurol.* 123, 3–16 (1992).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

84. Reier, P. J. , Stokes, B. T. , Thompson, F. J. & Anderson, P. J. Fetal cell grafts into resection injuries of the rat and cat spinal cord. *Exp. Neurol.* 115, 177–188 (1992).

- [Article](#)
- [PubMed](#)
- [ISI](#)

◦ **ChemPort**

85. Reier, P. J. , Anderson, D. K. , Young, W. , Michel, M. E. & Fessler, R. Workshop on intraspinal application. J. Neurotrauma 11, 369–377 (1994).

◦ **PubMed**

◦ **ChemPort**

86. Bregman, B. S. *et al.* Transplants and neurotrophic factors increase regeneration and recovery after spinal cord injury. Prog. Brain Res. 137, 257–273 (2002).

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

87. Coumans, J. V. *et al.* Axonal regeneration and functional recovery after complete spinal cord injury treated with transplants and neurotrophins. J. Neurosci. 21, 9334–9344 (2001).

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

88. Reier, P. J. Cellular transplantation strategies for spinal cord injury and translational neuroscience (2004).

Interesting review that describes in detail clinical trials that use cellular transplantation. Results from some trials have yet to be published in peer-reviewed articles.

◦ **Article**

◦ **PubMed**

89. Falcı, S. *et al.* Obliteration of a posttraumatic spinal cord cyst with solid human embryonic stem cells: a preliminary attempt. J. Neurotrauma 14, 875–884 (1997).

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

90. Thompson, F. J. *et al.* Neurophysiological assessment of the feasibility and safety of neurocyte transplantation in a rat model of syringomyelia. J. Neurotrauma 18, 931–945 (2001).

◦ **Article**

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

91. McDonald, J. W. *et al.* Transplanted embryonic stem cells survive, differentiate and promote neuroprotection in a rat model of spinal cord injury. Nature Med. 5, 1410–1412 (1999).

◦ **Article**

92. Harper, J. M. *et al.* Axonal growth of embryonic stem cell-derived motoneurons *in vitro* and *in vivo* in rats. Proc. Natl Acad. Sci. USA 101, 7123–7128 (2004).

◦ **Article**

◦ **PubMed**

◦ **ChemPort**

93. Cao, Q. L. , Howard, R. M. , Dennison, J. B. & Whittemore, S. R. Differentiation of engrafted embryonic stem cells into neurons after spinal cord injury in the rat. J. Neurosci. 24, 1100–1110 (2004).

cells is inhibited in the traumatically injured spinal cord. *Exp. Neurol.* 177, 349–359 (2002).

◦ [Article](#)

◦ [ChemPort](#)

94. Herrera, J. *et al.* Embryonic-derived glial-restricted precursor cells (GRP cells) can differentiate into oligodendrocytes *in vivo*. *Exp. Neurol.* 171, 11–21 (2001).

◦ [Article](#)

◦ [PubMed](#)

◦ [ChemPort](#)

95. Cao, Q. , Benton, R. L. & Whittemore, S. R. Stem cell repair of central nervous system injury. *Exp. Neurol.* 177, 360–375 (2002).

◦ [Article](#)

◦ [PubMed](#)

◦ [ChemPort](#)

96. Han, S. S. , Kang, D. Y. , Mujtaba, T. , Rao, M. S. & Fischer, I. Grafted lineage-restricted precursor cells differentiate into neurons in the adult spinal cord. *Exp. Neurol.* 177, 360–375 (2002).

◦ [Article](#)

97. Han, S. S. , Liu, Y. , Tyler-Polsz, C. , Rao, M. S. & Fischer, I. Transplantation of glial-restricted precursor cells into spinal cord: survival, glial-specific differentiation, and preferential migration in white matter. *Exp. Neurol.* 177, 360–375 (2002).

◦ [Article](#)

◦ [PubMed](#)

98. Yan, J. , Welsh, A. M. , Bora, S. H. , Snyder, E. Y. & Koliatsos, V. E. Differentiation and trophic support of neural precursors in the adult spinal cord. *J. Comp. Neurol.* 480, 101–114 (2004).

◦ [Article](#)

99. Hill, C. E. *et al.* Acute transplantation of glial-restricted precursor cells into spinal cord promotes axonal differentiation, and effects on lesion environment and axonal regeneration. *Exp. Neurol.* 177, 360–375 (2002).

◦ [Article](#)

◦ [PubMed](#)

◦ [ChemPort](#)

100. Ogawa, Y. *et al.* Transplantation of *in vitro*-expanded fetal neural progenitor cells results in functional recovery after spinal cord contusion injury in adult rats. *J. Neurosci. Res.* 69, 925–933 (2002).

◦ [Article](#)

◦ [PubMed](#)

◦ [ChemPort](#)

101. Teng, Y. D. *et al.* Functional recovery following traumatic spinal cord injury mediated by a novel population of endogenous neural stem cells. *Proc. Natl Acad. Sci. USA* 99, 3024–3029 (2002).

◦ [Article](#)

◦ [PubMed](#)

◦ [ChemPort](#)

102. Cummings, B. J. *et al.* Human neural stem cells differentiate and promote locomotor recovery after spinal cord injury. *Proc. Natl Acad. Sci. USA* 102, 14069–14074 (2005).

Important study in which transplanted cells are also ablated to determine whether recovered function.

- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
- 103. Iwanami, A. *et al.* Transplantation of human neural stem cells for spinal cord injury in pri
190 (2005).
- [Article](#)
 - [ChemPort](#)
- 104. Mitsui, T. , Shumsky, J. S. , Lepore, A. C. , Murray, M. & Fischer, I. Transplantation of neu
into contused spinal cord improves bladder and motor functions, decreases thermal hyper
circuitry. J. Neurosci. 25, 9624–9636 (2005).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
- 105. Keirstead, H. S. *et al.* Human embryonic stem cell-derived oligodendrocyte progenitor cel
restore locomotion after spinal cord injury. J. Neurosci. 25, 4694–4705 (2005).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 106. Faulkner, J. & Keirstead, H. S. Human embryonic stem cell-derived oligodendrocyte prog
cord injury. Transpl. Immunol. 15, 131–142 (2005).
- [ChemPort](#)
- 107. Schultz, S. S. Adult stem cell application in spinal cord injury. Curr. Drug Targets 6, 63–73
(2005).
- [Article](#)
 - [ChemPort](#)
- 108. Koshizuka, S. *et al.* Transplanted hematopoietic stem cells from bone marrow differentiat
promote functional recovery after spinal cord injury in mice. J. Neuropathol. Exp. Neurol.
157, 105–115 (2005).
- 109. Koda, M. *et al.* Hematopoietic stem cell and marrow stromal cell for spinal cord injury in
(2005).
- [Article](#)
- 110. Hofstetter, C. P. *et al.* Allodynia limits the usefulness of intraspinal neural stem cell grafts
outcome. Nature Neurosci. 8, 346–353 (2005).
- [Article](#)
- 111. Wu, S. *et al.* Bone marrow stromal cells enhance differentiation of cocultured neurosphere
injured spinal cord. J. Neurosci. Res. 72, 343–351 (2003).
- [Article](#)
 - [ChemPort](#)

112. Reynolds, B. A. & Weiss, S. Generation of neurons and astrocytes from isolated cells of the system. *Science* 255, 1707–1710 (1992).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
113. Gage, F. H. *et al.* Survival and differentiation of adult neuronal progenitor cells transplant Acad. Sci. USA 92, 11879–11883 (1995).
 - [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
114. Shihabuddin, L. S. , Horner, P. J. , Ray, J. & Gage, F. H. Adult spinal cord stem cells gener the adult dentate gyrus. *J. Neurosci.* 20, 8727–8735 (2000).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
115. Vroemen, M. , Aigner, L. , Winkler, J. & Weidner, N. Adult neural progenitor cell grafts su and integrate along axonal pathways. *Eur. J. Neurosci.* 18, 743–751 (2003).
 - [Article](#)
 - [PubMed](#)
116. Cao, Q. L. *et al.* Pluripotent stem cells engrafted into the normal or lesioned adult rat spin lineage. *Exp. Neurol.* 167, 48–58 (2001).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
117. Karimi-Abdolrezaee, S. , Eftekharpour, E. , Wang, J. , Morshead, C. M. & Fehlings, M. G. I neural precursor cells promotes remyelination and functional neurological recovery after : 3377–3389 (2006).
 - [Article](#)
 - [ChemPort](#)
118. Horner, P. J. *et al.* Proliferation and differentiation of progenitor cells throughout the inta 20, 2218–2228 (2000).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
119. Howard, M. J. , Liu, S. , Schottler, F. , Snider, B. J. & Jacquin, M. F. Transplantation of ap cells into the injured rat spinal cord. *Somatosens. Mot. Res.* 22, 37–44 (2005).
 - [Article](#)
120. Chen, J. , Bernreuther, C. , Dihne, M. & Schachner, M. Cell adhesion molecule L1-transfec enhanced survival support regrowth of corticospinal tract axons in mice after spinal cord i (2005).

- [Article](#)
- 121. Cao, Q. *et al.* Functional recovery in traumatic spinal cord injury after transplantation of restricted precursor cells. *J. Neurosci.* 25, 6947–6957 (2005).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 122. Hamada, M. *et al.* Introduction of the MASH1 gene into mouse embryonic stem cells leads to precursors lacking Nogo receptor expression that can be applicable for transplantation to CNS. *J. Neurosci.* 22, 509–522 (2006).
- [Article](#)
 - [ChemPort](#)
- 123. Hirschberg, D. L. & Schwartz, M. Macrophage recruitment to acutely injured central nervous system: a basis for an immune-brain barrier. *J. Neuroimmunol.* 61, 89–96 (1995).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 124. Rapalino, O. *et al.* Implantation of stimulated homologous macrophages results in partial recovery of locomotor function after spinal cord injury. *Neurosci. Lett.* 4, 814–821 (1998).
- [Article](#)
- 125. Bomstein, Y. *et al.* Features of skin-coincubated macrophages that promote recovery from spinal cord injury. *J. Neuroimmunol.* 142, 10–16 (2003).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 126. Moon, L. & Bunge, M. B. From animal models to humans: strategies for promoting CNS axon growth and limb function after spinal cord injury. *J. Neurol. Phys. Ther.* 29, 55–69 (2005).
- 127. Popovich, P. G. *et al.* The neuropathological and behavioral consequences of intraspinal injection of macrophages after spinal cord injury. *Neurosci. Lett.* 61, 623–633 (2002).
- [ChemPort](#)
- 128. Popovich, P. G. *et al.* Depletion of hematogenous macrophages promotes partial hindlimb repair after experimental spinal cord injury. *Exp. Neurol.* 158, 351–365 (1999).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
- 129. Knoller, N. *et al.* Clinical experience using incubated autologous macrophages as a treatment for spinal cord injury: Phase I study results. *J. Neurosurg. Spine* 3, 173–181 (2005).

130. Baptiste, D. C. & Fehlings, M. G. Pharmacological approaches to repair the injured spinal cord. *Neurotrauma* 20, 1017–1027 (2003).
 - [Article](#)
131. Gris, D. *et al.* Transient blockade of the CD11d/CD18 integrin reduces secondary damage to sensory, autonomic, and motor function. *J. Neurosci.* 24, 4043–4051 (2004).
 - [Article](#)
 - [ChemPort](#)
132. Lee, S. M. *et al.* Minocycline reduces cell death and improves functional recovery after traumatic spinal cord injury. *J. Neurotrauma* 20, 1017–1027 (2003).
 - [Article](#)
133. Teng, Y. D. *et al.* Minocycline inhibits contusion-triggered mitochondrial cytochrome c release and deficits after spinal cord injury. *Proc. Natl Acad. Sci. USA* 101, 3071–3076 (2004).
 - [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
134. Stirling, D. P. *et al.* Minocycline treatment reduces delayed oligodendrocyte death, attenuates axonal loss, and improves functional outcome after spinal cord injury. *J. Neurosci.* 24, 2182–2190 (2004).
 - [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
135. Hall, E. D. & Springer, J. E. Neuroprotection and acute spinal cord injury: a reappraisal. *Neurotrauma* 20, 1017–1027 (2003).
 - [Article](#)
136. Coleman, W. P. *et al.* A critical appraisal of the reporting of the National Acute Spinal Cord Injury Study. *J. Spinal Disord.* 13, 185–199 (2000).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
137. Hurlbert, R. J. Methylprednisolone for acute spinal cord injury: an inappropriate standard. *Neurotrauma* 20, 1017–1027 (2003).
 - [PubMed](#)
 - [ChemPort](#)
138. Fehlings, M. G. & Baptiste, D. C. Current status of clinical trials for acute spinal cord injury. *Neurotrauma* 20, 1017–1027 (2003).
 - [Article](#)
139. Short, D. J. , El Masry, W. S. & Jones, P. W. High dose methylprednisolone in the management of acute spinal cord injury: a systematic review from a clinical perspective. *Spinal Cord* 38, 273–286 (2000).
 - [Article](#)
 - [PubMed](#)

- **ChemPort**
- 140. Bracken, M. B. Methylprednisolone and acute spinal cord injury: an update of the random (2001).
- **Article**
 - **ChemPort**
- 141. McTigue, D. M. , Horner, P. J. , Stokes, B. T. & Gage, F. H. Neurotrophin-3 and brain-deri oligodendrocyte proliferation and myelination of regenerating axons in the contused adult 5354–5365 (1998).
- **PubMed**
 - **ISI**
 - **ChemPort**
- 142. Houweling, D. A. *et al.* Local application of collagen containing brain-derived neurotrophin function after spinal cord injury in the adult rat. *Neurosci. Lett.* 251, 193–196 (1998).
- **Article**
 - **PubMed**
 - **ChemPort**
- 143. Jakeman, L. B. , Wei, P. , Guan, Z. & Stokes, B. T. Brain-derived neurotrophic factor stimu sprouting of cholinergic fibers after spinal cord injury. *Exp. Neurol.* 154, 170–184 (1998).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 144. Namiki, J. , Kojima, A. & Tator, C. H. Effect of brain-derived neurotrophic factor, nerve gr functional recovery and regeneration after spinal cord injury in adult rats. *J. Neurotrauma*
- **PubMed**
 - **ISI**
 - **ChemPort**
- 145. Shumsky, J. S. *et al.* Delayed transplantation of fibroblasts genetically modified to secrete injury site is associated with limited recovery of function. *Exp. Neurol.* 184, 114–130 (2003).
- **Article**
 - **ChemPort**
- 146. Tobias, C. A. *et al.* Delayed grafting of BDNF and NT-3 producing fibroblasts into the inju sprouting, partially rescues axotomized red nucleus neurons from loss and atrophy, and p *Neurol.* 184, 97–113 (2003).
- **Article**
 - **ChemPort**
- 147. Zhou, L. & Shine, H. D. Neurotrophic factors expressed in both cortex and spinal cord ind cord injury. *J. Neurosci. Res.* 74, 221–226 (2003).
- **Article**
 - **ChemPort**
- 148. Lee, T. T. , Green, B. A. , Dietrich, W. D. & Yeziarski, R. P. Neuroprotective effects of basic

spinal cord contusion injury in the rat. *J. Neurotrauma* 16, 347–356 (1999).

◦ **ChemPort**

149. Blesch, A. & Tuszynski, M. H. Cellular GDNF delivery promotes growth of motor and dorsal partial and complete spinal cord transections and induces remyelination. *J. Comp. Neurol.*

◦ **Article**

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

150. Tuszynski, M. H. , Grill, R. , Jones, L. L. , McKay, H. M. & Blesch, A. Spontaneous and augmented primate spinal cord: effects of local injury and nerve growth factor-secreting cell grafts. *J.*

◦ **Article**

◦ **ChemPort**

151. Tuszynski, M. H. *et al.* NT-3 gene delivery elicits growth of chronically injured corticospinal functional deficits after chronic scar resection. *Exp. Neurol.* 181, 47–56 (2003).

◦ **Article**

◦ **ChemPort**

152. Mitsui, T. , Fischer, I. , Shumsky, J. S. & Murray, M. Transplants of fibroblasts expressing of bladder and hindlimb function following spinal contusion injury in rats. *Exp. Neurol.* 195

◦ **Article**

◦ **ChemPort**

153. Blesch, A. , Yang, H. , Weidner, N. , Hoang, A. & Otero, D. Axonal responses to cellularly derived injury. *Mol. Cell. Neurosci.* 27, 190–201 (2004).

◦ **Article**

◦ **ChemPort**

154. Schnell, L. , Schneider, R. , Kolbeck, R. , Barde, Y. A. & Schwab, M. E. Neurotrophin-3 enhances tract during development and after adult spinal cord lesions. *Nature* 367, 170–172 (1994).

◦ **Article**

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

155. Zhou, L. , Baumgartner, B. J. , Hill-Felberg, S. J. , McGowen, L. R. & Shine, H. D. Neurotrophin-3 axonal plasticity in the adult injured spinal cord. *J. Neurosci.* 23, 1424–1431 (2003).

◦ **ChemPort**

156. Koda, M. *et al.* Adenovirus vector-mediated *in vivo* gene transfer of brain-derived neurotrophic factor promotes rubrospinal axonal regeneration and functional recovery after complete transection of the spinal cord. *Neurotrauma* 21, 329–337 (2004).

◦ **Article**

157. Blits, B. & Bunge, M. B. Direct gene therapy for repair of the spinal cord. *J. Neurotrauma* : **Thorough review of viral vectors that have been used to deliver molecules of interest**

- **Article**
- 158. Grill, R. , Blesch, A. & Tuszynski, M. H. Robust growth of chronically injured spinal cord a modified NGF-secreting cells. *Exp. Neurol.* 148, 444–452 (1997).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 159. Tuszynski, M. H. *et al.* Nerve growth factor delivery by gene transfer induces differential o noradrenergic neurites after adult spinal cord injury. *Exp. Neurol.* 137, 157–173 (1996).
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 160. Grill, R. , Murai, K. , Blesch, A. , Gage, F. H. & Tuszynski, M. H. Cellular delivery of neurot axon growth and partial functional recovery after spinal cord injury. *J. Neurosci.* 17, 5560.
- **PubMed**
 - **ISI**
 - **ChemPort**
- 161. Jin, Y. , Fischer, I. , Tessler, A. & Houle, J. D. Transplants of fibroblasts genetically modifi regeneration from supraspinal neurons following chronic spinal cord injury. *Exp. Neurol.*
- **Article**
 - **PubMed**
 - **ISI**
 - **ChemPort**
- 162. Jin, Y. , Tessler, A. , Fischer, I. & Houle, J. D. Fibroblasts genetically modified to produce : chronically injured serotonergic axons. *Neurorehabil. Neural Repair* 14, 311–317 (2000).
- **ChemPort**
- 163. Liu, Y. *et al.* Transplants of fibroblasts genetically modified to express BDNF promote reg axons and recovery of forelimb function. *J. Neurosci.* 19, 4370–4387 (1999).
- **PubMed**
 - **ISI**
 - **ChemPort**
- 164. Lu, P. , Blesch, A. & Tuszynski, M. H. Neurotrophism without neurotropism: BDNF promc lesioned corticospinal neurons. *J. Comp. Neurol.* 436, 456–470 (2001).
- **Article**
 - **ChemPort**
- 165. Lu, P. , Jones, L. L. & Tuszynski, M. H. BDNF-expressing marrow stromal cells support ex spinal cord injury. *Exp. Neurol.* 191, 344–360 (2005).
- **Article**
 - **PubMed**
 - **ChemPort**

166. Nothias, J. M. *et al.* Combined effects of neurotrophin secreting transplants, exercise, and improve function in spinal rats. *Neurorehabil. Neural Repair* 19, 296–312 (2005).
 - [Article](#)
167. Kwon, B. K. *et al.* Rubrospinal neurons fail to respond to brain-derived neurotrophic factor 2 months after cervical axotomy. *Exp. Neurol.* 189, 45–57 (2004).
 - [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
168. Apfel, S. C. Nerve growth factor for the treatment of diabetic neuropathy: what went wrong in the future hold? *Int. Rev. Neurobiol.* 50, 393–413 (2002).
 - [PubMed](#)
 - [ChemPort](#)
169. Tuszynski, M. H. *et al.* A phase 1 clinical trial of nerve growth factor gene therapy for Alzheimer's disease. *Neuron* 47, 555 (2005).
 - [Article](#)
170. Cai, D. *et al.* Neuronal cyclic AMP controls the developmental loss in ability of axons to regenerate. *Neuron* 30, 101–112 (2001).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
171. Neumann, S. , Bradke, F. , Tessier-Lavigne, M. & Basbaum, A. I. Regeneration of sensory neurons is induced by intraganglionic cAMP elevation. *Neuron* 34, 885–893 (2002).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
172. Kao, H. T. *et al.* A protein kinase A-dependent molecular switch in synapsins regulates neuronal function. *Neuron* 34, 431–437 (2002).
 - [Article](#)
173. Qiu, J. *et al.* Spinal axon regeneration induced by elevation of cyclic AMP. *Neuron* 34, 895–905 (2002).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
174. Lu, P. , Yang, H. , Jones, L. L. , Filbin, M. T. & Tuszynski, M. H. Combinatorial therapy with neurotrophins and neurotrophin receptors promotes axonal regeneration beyond sites of spinal cord injury. *J. Neurosci.* 24, 6402–6412 (2004).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)

175. Bhatt, D. H. , Otto, S. J. , Depoister, B. & Fetcho, J. R. Cyclic AMP-induced repair of zebra
254–258 (2004).
 - o [Article](#)
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
176. Nikulina, E. , Tidwell, J. L. , Dai, H. N. , Bregman, B. S. & Filbin, M. T. The phosphodiesterase
after a spinal cord lesion promotes axonal regeneration and functional recovery. *Proc. Natl. Acad. Sci. USA* 101, 1511–1516 (2004).
 - o [Article](#)
 - o [PubMed](#)
 - o [ChemPort](#)
177. Pearse, D. D. *et al.* cAMP and Schwann cells promote axonal growth and functional recovery after
spinal cord injury. *Neuron* 40, 610–616 (2004).
 - o [Article](#)
178. Niederost, B. , Oertle, T. , Fritsche, J. , McKinney, R. A. & Bandtlow, C. E. Nogo-A and myelin
neurite growth inhibition by antagonistic regulation of RhoA and Rac1. *J. Neurosci.* 22, 1011–1021 (2002).
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
179. Jain, A. , Brady-Kalnay, S. M. & Bellamkonda, R. V. Modulation of Rho GTPase activity alleviates
proteoglycan-dependent inhibition of neurite extension. *J. Neurosci. Res.* 77, 299–307 (2004).
 - o [Article](#)
 - o [PubMed](#)
 - o [ChemPort](#)
180. Schweigreiter, R. *et al.* Versican V2 and the central inhibitory domain of Nogo-A inhibit neurite
extension through independent pathways that converge at RhoA. *Mol. Cell. Neurosci.* 27, 163–174 (2004).
 - o [Article](#)
 - o [PubMed](#)
 - o [ChemPort](#)
181. Dergham, P. *et al.* Rho signaling pathway targeted to promote spinal cord repair. *J. Neurosci.* 23, 1416–1423 (2003).
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
182. Fournier, A. E. , Takizawa, B. T. & Strittmatter, S. M. Rho kinase inhibition enhances axonal
regeneration after spinal cord injury. *J. Neurosci.* 23, 1416–1423 (2003).
 - o [PubMed](#)
 - o [ISI](#)
 - o [ChemPort](#)
183. Sung, J. K. *et al.* A possible role of RhoA/Rho-kinase in experimental spinal cord injury in
rodents. *Neurosci. Lett.* 344, 141–144 (2003).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

184. McKerracher, L. & Higuchi, H. Targeting Rho to stimulate repair after spinal cord injury. (2006).

- [Article](#)

185. Chan, C. C. *et al.* Dose-dependent beneficial and detrimental effects of ROCK inhibitor Y27 on functional recovery after rat spinal cord injury. *Exp. Neurol.* 196, 352–364 (2005).

- [Article](#)
- [ChemPort](#)

186. Mueller, B. K., Mack, H. & Teusch, N. Rho kinase, a promising drug target for neurological disorders. *Discov.* 4, 387–398 (2005).

Good review on the potential of ROCK inhibitors to prevent neurodegeneration and promote neuroregeneration in various neurological disorders, including SCI.

- [Article](#)

187. Hara, M. *et al.* Protein kinase inhibition by fasudil hydrochloride promotes neurological recovery in rats. *J. Neurosurg.* 93, 94–101 (2000).

- [ChemPort](#)

188. Tanaka, H. *et al.* Cytoplasmic p21(Cip1/WAF1) enhances axonal regeneration and function in rats. *Neuroscience* 127, 155–164 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

189. Wei, L. *et al.* Rho kinases play an obligatory role in vertebrate embryonic organogenesis. *J. Neurosci.* 21, 1000–1008 (2001).

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

190. Niederöst, B. P., Zimmermann, D. R., Schwab, M. E. & Bandtlow, C. E. Bovine CNS myelin inhibitory activity associated with chondroitin sulfate proteoglycans. *J. Neurosci.* 19, 8979–8988 (1999).

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

191. Schwab, M. E. Nogo and axon regeneration. *Curr. Opin. Neurobiol.* 14, 118–124 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

192. Fouad, K., Klusman, I. & Schwab, M. E. Regenerating corticospinal fibers in the Marmoset monkey. *J. Neurosci.* 19, 8979–8988 (1999).

cord lesion and treatment with the anti-Nogo-A antibody IN-1. *Eur. J. Neurosci.* 20, 2479.

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

193. Benson, M. D. *et al.* Ephrin-B3 is a myelin-based inhibitor of neurite outgrowth. *Proc. Natl* (2005).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

194. Schnell, L. & Schwab, M. E. Axonal regeneration in the rat spinal cord produced by an ant neurite growth inhibitors. *Nature* 343, 269–272 (1990).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

195. Schnell, L. & Schwab, M. E. Sprouting and regeneration of lesioned corticospinal tract fibre *J. Neurosci.* 5, 1156–1171 (1993).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

196. Bregman, B. S. *et al.* Recovery from spinal cord injury by antibodies to neurite growth inh (1995).

- [Article](#)
- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

197. von Meyenburg, J. , Brosamle, C. , Metz, G. A. & Schwab, M. E. Regeneration and sproutin corticospinal tract fibers in adult rats promoted by NT-3 and the mAb IN-1, which neutral growth inhibitors. *Exp. Neurol.* 154, 583–594 (1998).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

198. Thallmair, M. *et al.* Neurite growth inhibitors restrict plasticity and functional recovery fo *Nature Neurosci.* 1, 124–131 (1998).

- [Article](#)

199. Z'Graggen, W. J. , Metz, G. A. S. , Kaitje, G. L. , Thallmair, M. & Schwab, M. E. Functional plasticity after unilateral pyramidal tract lesion and blockade of myelin-associated neurite *Neurosci.* 18, 4744–4757 (1998).

- [PubMed](#)
- [ChemPort](#)

200. Oudega, M. *et al.* Neutralizing antibodies against neurite growth inhibitor ni-35/250 do not promote axons in the adult rat spinal cord. *Neuroscience* 100, 873–883 (2000).
 - [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
201. Guest, J. D. *et al.* Influence of IN-1 antibody and acidic FGF-fibrin glue on the response of human Schwann cell grafts. *J. Neurosci. Res.* 50, 888–905 (1997).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
202. Merkler, D. *et al.* Locomotor recovery in spinal cord-injured rats treated with an antibody against neurite growth inhibitor Nogo-A. *J. Neurosci.* 21, 3665–3673 (2001).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
203. Mandemakers, W. J. & Barres, B. A. Axon regeneration: it's getting crowded at the gates of the spinal cord (2005).
 - [Article](#)
 - [ChemPort](#)
204. Song, X. Y. , Zhong, J. H. , Wang, X. & Zhou, X. F. Suppression of p75^{NTR} does not promote axon regeneration in the spinal cord in mice. *J. Neurosci.* 24, 542–546 (2004).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
205. Kim, J. E. , Li, S. , GrandPre, T. , Qiu, D. & Strittmatter, S. M. Axon regeneration in young mice after spinal cord injury. *Neuron* 38, 187–199 (2003).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
206. Simonen, M. *et al.* Systemic deletion of the myelin-associated outgrowth inhibitor Nogo-A promotes axon regeneration responses after spinal cord injury. *Neuron* 38, 201–211 (2003).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
207. Zheng, B. *et al.* Lack of enhanced spinal regeneration in Nogo-deficient mice. *Neuron* 38, 101–113 (2002).
 - [Article](#)
 - [PubMed](#)

- [ISI](#)
 - [ChemPort](#)
208. Zheng, B. *et al.* Genetic deletion of the Nogo receptor does not reduce neurite inhibition in tract regeneration *in vivo*. *Proc. Natl Acad. Sci. USA* 102, 1205–1210 (2005).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
209. Li, S. *et al.* Blockade of Nogo-66, myelin-associated glycoprotein, and oligodendrocyte myelin glycoprotein promotes axonal sprouting and recovery after spinal injury. *J. Neurosci.* 24, 1066–1074 (2004).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
210. Ji, B. *et al.* Effect of combined treatment with methylprednisolone and soluble Nogo-66 receptor on axonal sprouting and recovery after spinal cord injury. *Eur. J. Neurosci.* 22, 587–594 (2005).
- [Article](#)
211. Li, S. & Strittmatter, S. M. Delayed systemic Nogo-66 receptor antagonist promotes recovery after spinal cord injury. *J. Neurosci.* 23, 4219–4227 (2003).
- [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
212. GrandPre, T., Li, S. & Strittmatter, S. M. Nogo-66 receptor antagonist peptide promotes axonal sprouting and recovery after spinal cord injury. *J. Neurosci.* 22, 547–551 (2002).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
213. Stichel, C. C. *et al.* Inhibition of collagen IV deposition promotes regeneration of CNS axons after spinal cord injury. *J. Neurosci.* 19, 1000–1009 (1999).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
214. Klapka, N. *et al.* Suppression of fibrous scarring in spinal cord injury of rat promotes long corticospinal tract axons, rescue of primary motoneurons in somatosensory cortex and improves locomotor recovery. *J. Neurosci.* 22, 3047–3058 (2005).
- [Article](#)
215. Weidner, N., Grill, R. J. & Tuszynski, M. H. Elimination of basal lamina and the collagenous scar augment corticospinal tract regeneration. *Exp. Neurol.* 160, 40–50 (1999).
- [Article](#)
 - [ChemPort](#)

216. Moon, L. D. , Asher, R. A. , Rhodes, K. E. & Fawcett, J. W. Regeneration of CNS axons back of adult rat brain with chondroitinase ABC. *Nature Neurosci.* 4, 465–466 (2001).
217. Bradbury, E. J. *et al.* Chondroitinase ABC promotes functional recovery after spinal cord injury.
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
218. Yick, L. W. , Wu, W. , So, K. F. , Yip, H. K. & Shum, D. K. Chondroitinase ABC promotes axonal growth of neurons after spinal cord injury. *Neuroreport* 11, 1063–1067 (2000).
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
219. Chau, C. H. *et al.* Chondroitinase ABC enhances axonal regrowth through Schwann cell-seeded scaffolds after spinal cord injury. *FASEB J.* 18, 194–196 (2004).
 - [PubMed](#)
 - [ChemPort](#)
220. Fouad, K. *et al.* Combining Schwann cell bridges and olfactory-ensheathing glia grafts with neurotrophins promotes locomotor recovery after complete transection of the spinal cord. *J. Neurosci.* 25, 1169–1177 (2005).
 - [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
221. Caggiano, A. O. , Zimmer, M. P. , Ganguly, A. , Blight, A. R. & Gruskin, E. A. Chondroitinase ABC improves bladder function following contusion injury of the rat spinal cord. *J. Neurotrauma* 22, 226–236 (2005).
 - [Article](#)
 - [PubMed](#)
222. Engesser-Cesar, C. , Anderson, A. J. , Basso, D. M. , Edgerton, V. R. & Cotman, C. W. Voluntary recovery from a moderate spinal cord injury. *J. Neurotrauma* 22, 157–171 (2005).
 - [Article](#)
 - [PubMed](#)
223. Ichiyama, R. M. , Gerasimenko, Y. P. , Zhong, H. , Roy, R. R. & Edgerton, V. R. Hindlimb recovery in spinal rats induced by epidural spinal cord stimulation. *Neurosci. Lett.* 383, 339–344 (2005).
 - [Article](#)
 - [ChemPort](#)
224. Bouyer, L. J. Animal models for studying potential training strategies in persons with spinal cord injury. *Ther. Adv. Neurol.* 29, 117–125 (2005).
225. Curt, A. , Schwab, M. E. & Dietz, V. Providing the clinical basis for new interventional therapies: the assessment of recovery after spinal cord injury. *Spinal Cord* 42, 1–6 (2004).
 - [Article](#)

- PubMed
- ISI
- ChemPort

226. Edgerton, V. R. , Kim, S. J. , Ichiyama, R. M. , Gerasimenko, Y. P. & Roy, R. R. Rehabilitation after spinal cord injury. *J. Neurotrauma* 23, 560–570 (2006).

Excellent review on locomotor rehabilitation after SCI that includes a comprehensive review of automaticity and plasticity of the spinal circuitry.

- Article

227. Edgerton, V. R. , Tillakaratne, N. J. , Bigbee, A. J. , de Leon, R. D. & Roy, R. R. Plasticity of the spinal cord after injury. *Annu. Rev. Neurosci.* 27, 145–167 (2004).

- Article
- PubMed
- ChemPort

228. Moon, L. D. F. , Leasure, J. L. , Gage, F. H. & Bunge, M. B. Environmental enrichment sustains functional recovery following spinal cord transection and glial transplantation. *Restor. Neurol. Neurosci.* (in the Press).

229. Lankhorst, A. J. *et al.* Effects of enriched housing on functional recovery after spinal cord injury. *J. Neurotrauma* 18, 203–215 (2001).

- Article
- ChemPort

230. Hutchinson, K. J. , Gomez-Pinilla, F. , Crowe, M. J. , Ying, Z. & Basso, D. M. Three exercises promote sensory recovery after spinal cord contusion in rats. *Brain* 127, 1403–1414 (2004).

- Article

231. Van Meeteren, N. L. , Eggers, R. , Lankhorst, A. J. , Gispén, W. H. & Hamers, F. P. Locomotor recovery after spinal cord contusion injury in rats is improved by spontaneous exercise. *J. Neurotrauma* 20, 1029–1039 (2003).

- Article

232. Barbeau, H. , Nadeau, S. & Garneau, C. Physical determinants, emerging concepts, and treatment approaches for individuals with spinal cord injury. *J. Neurotrauma* 23, 571–585 (2006).

- Article

233. Field-Fote, E. C. , Lindley, S. D. & Sherman, A. L. Locomotor training approaches for individuals with spinal cord injury: preliminary report of walking-related outcomes. *J. Neurol. Phys. Ther.* 29, 127–137 (2005).

234. Wolpaw, J. R. Treadmill training after spinal cord injury: good but not better. *Neurology* 64, 1000–1001 (2005).

- Article
- PubMed

235. Dobkin, B. *et al.* Weight-supported treadmill vs over-ground training for walking after acute spinal cord injury. *J. Neurotrauma* 23, 484–493 (2006).

- Article
- PubMed
- ChemPort

236. Herman, R. , He, J. , D'Luzansky, S. , Willis, W. & Dilli, S. Spinal cord stimulation facilitates recovery of locomotion after spinal cord injury. *J. Neurotrauma* 23, 1000–1001 (2006).

incomplete spinal cord injured. *Spinal Cord* 40, 65–68 (2002).

- [Article](#)
- [ChemPort](#)

237. Wernig, A. Long-term body-weight supported treadmill training and subsequent follow-up effects on functional walking ability and measures of subjective well-being. *Spinal Cord* 40, 65–68 (2002).

- [Article](#)

238. Humm, J. L. , Kozlowski, D. A. , James, D. C. , Gotts, J. E. & Schallert, T. Use-dependent effects during an early post-lesion vulnerable period. *Brain Res.* 783, 286–292 (1998).

- [Article](#)
- [PubMed](#)
- [ChemPort](#)

239. de Leon, R. D. , Hodgson, J. A. , Roy, R. R. & Edgerton, V. R. Locomotor capacity attributes spontaneous recovery after spinalization in adult cats. *J. Neurophysiol.* 79, 1329–1340 (1998).

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

240. Lu, J. & Ashwell, K. Olfactory ensheathing cells: their potential use for repairing the injured spinal cord. *Spinal Cord* 40, 65–68 (2002).

- [Article](#)

241. Morrissey, T. K. , Kleitman, N. & Bunge, R. P. Isolation and functional characteristics of Schwann cells from peripheral nerve. *J. Neurosci.* 11, 2433–2442 (1991).

- [PubMed](#)
- [ISI](#)
- [ChemPort](#)

242. Emery, E. , Li, X. , Brunschwig, J. P. , Olson, L. & Levi, A. D. Assessment of the malignant potential of human Schwann cells. *J. Peripher. Nerv. Syst.* 4, 107–116 (1999).

- [ChemPort](#)

243. Lepore, A. C. , Bakshi, A. , Swanger, S. A. , Rao, M. S. & Fischer, I. Neural precursor cells migrate to the cervical spinal cord by intrathecal injection at the lumbar cord. *Brain Res.* 1045, 206–216 (2000).

- [ChemPort](#)

244. Bakshi, A. *et al.* Lumbar puncture delivery of bone marrow stromal cells in spinal cord contusion: a minimally invasive cell transplantation. *J. Neurotrauma* 23, 55–65 (2006).

- [Article](#)

245. Ramer, L. M. *et al.* Peripheral olfactory ensheathing cells reduce scar and cavity formation after spinal cord injury. *J. Comp. Neurol.* 473, 1–15 (2004).

- [Article](#)
- [PubMed](#)
- [ISI](#)

246. Hill, C. E. , Moon, L. D. , Wood, P. M. & Bunge, M. B. Labeled Schwann cell transplantation in the injured spinal cord. *Spinal Cord* 40, 65–68 (2002).

replacement, and strategies to enhance survival. *Glia* 53, 338–343 (2006).

◦ **Article**

247. Barakat, D. J. *et al.* Survival, integration, and axon growth support of glia transplanted into cord. *Cell Transplant.* 14, 225–240 (2005).

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

248. Hendriks, W. T. , Ruitenberg, M. J. , Blits, B. , Boer, G. J. & Verhaagen, J. Viral vector-mediated neurotrophins to promote regeneration of the injured spinal cord. *Prog. Brain Res.* 146, 41–50 (2005).

◦ **Article**

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

249. Kaspar, B. K. *et al.* Adeno-associated virus effectively mediates conditional gene modification in the CNS. *Sci. USA* 99, 2320–2325 (2002).

◦ **Article**

◦ **PubMed**

◦ **ChemPort**

250. Pirozzi, M. *et al.* Intramuscular viral delivery of paracalpain rescues peripheral axonopathy in paraplegia. *J. Clin. Invest.* 116, 202–208 (2006).

◦ **ChemPort**

251. Bresnahan, J. C. An electron-microscopic analysis of axonal alterations following blunt contusion of the thoracic spinal cord in the rhesus monkey (*Macaca mulatta*). *J. Neurol. Sci.* 37, 59–82 (1978).

◦ **Article**

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

252. Bresnahan, J. C. , King, J. S. , Martin, G. F. & Yashon, D. A neuroanatomical analysis of spinal cord injury in the rhesus monkey (*Macaca mulatta*). *J. Neurol. Sci.* 28, 521–542 (1976).

◦ **Article**

◦ **PubMed**

◦ **ISI**

◦ **ChemPort**

253. Courtine, G. *et al.* Performance of locomotion and foot grasping following a unilateral thoracic spinal cord injury in rhesus monkeys (*Macaca mulatta*). *Brain* 128, 2338–2358 (2005).

◦ **Article**

254. Hagg, T. & Oudega, M. Degenerative and spontaneous regenerative processes after spinal cord injury. *Neurosci. Biobehav. Rev.* 30, 263–280 (2006).

◦ **Article**

255. Richter, M. W. , Fletcher, P. A. , Liu, J. , Tetzlaff, W. & Roskams, A. J. Lamina propria and

exhibit differential integration and migration and promote differential axon sprouting in t
25, 10700–10711 (2005).

◦ [Article](#)

◦ [ChemPort](#)

256. Radojicic, M. , Reier, P. J. , Steward, O. & Keirstead, H. S. Septations in chronic spinal cor
Neurol. 196, 339–341 (2005).

◦ [Article](#)

257. Ye, J. H. & Houle, J. D. Treatment of the chronically injured spinal cord with neurotrophic
regeneration from supraspinal neurons. Exp. Neurol. 143, 70–81 (1997).

◦ [Article](#)

◦ [PubMed](#)

◦ [ISI](#)

◦ [ChemPort](#)

258. Kwon, B. K. *et al.* Survival and regeneration of rubrospinal neurons 1 year after spinal cor
99, 3246–3251 (2002).

◦ [Article](#)

◦ [PubMed](#)

◦ [ChemPort](#)

259. Pittenger, M. F. *et al.* Multilineage potential of adult human mesenchymal stem cells. Scie

◦ [Article](#)

◦ [PubMed](#)

◦ [ISI](#)

◦ [ChemPort](#)

260. Punzel, M. & Ho, A. D. Divisional history and pluripotency of human hematopoietic stem
(2001).

◦ [ChemPort](#)

261. Lee, J. *et al.* Migration and differentiation of nuclear fluorescence-labeled bone marrow st
into cerebral infarct and spinal cord injury in mice. Neuropathology 23, 169–180 (2003).

◦ [Article](#)

◦ [PubMed](#)

262. Woodbury, D. , Schwarz, E. J. , Prockop, D. J. & Black, I. B. Adult rat and human bone ma
neurons. J. Neurosci. Res. 61, 364–370 (2000).

◦ [Article](#)

◦ [PubMed](#)

◦ [ISI](#)

◦ [ChemPort](#)

263. Mezey, E. , Chandross, K. J. , Harta, G. , Maki, R. A. & McKercher, S. R. Turning blood into
antigens generated *in vivo* from bone marrow. Science 290, 1779–1782 (2000).

◦ [Article](#)

◦ [PubMed](#)

- [ISI](#)
 - [ChemPort](#)
264. Brazelton, T. R. , Rossi, F. M. , Keshet, G. I. & Blau, H. M. From marrow to brain: expressi mice. *Science* 290, 1775–1779 (2000).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
265. Castro, R. F. *et al.* Failure of bone marrow cells to transdifferentiate into neural cells *in vi*
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
266. Wurmser, A. E. & Gage, F. H. Stem cells: cell fusion causes confusion. *Nature* 416, 485–48
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
267. Terada, N. *et al.* Bone marrow cells adopt the phenotype of other cells by spontaneous cell (2002).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
268. Camargo, F. D. , Chambers, S. M. & Goodell, M. A. Stem cell plasticity: from transdifferent Prolif. 37, 55–65 (2004).
- [Article](#)
 - [PubMed](#)
 - [ISI](#)
 - [ChemPort](#)
269. Lu, P. & Tuszynski, M. H. Can bone marrow-derived stem cells differentiate into functiona 278 (2005).
- [Article](#)
 - [ChemPort](#)
270. Hofstetter, C. P. *et al.* Marrow stromal cells form guiding strands in the injured spinal cor Acad. Sci. USA 99, 2199–2204 (2002).
- [Article](#)
 - [PubMed](#)
 - [ChemPort](#)
271. Svendsen, C. N. , Bhattacharyya, A. & Tai, Y. T. Neurons from stem cells: preventing an id 2, 831–834 (2001).

- **Article**

272. Cogle, C. R. *et al.* Bone marrow transdifferentiation in brain after transplantation: a retrospective study. *Stem Cells* 24, 1437 (2004).

- **Article**

- **PubMed**

- **ISI**

- **ChemPort**

273. Deng, J. , Petersen, B. E. , Steindler, D. A. , Jorgensen, M. L. & Laywell, E. D. Mesenchymal stem cells express neural proteins in culture, and are neurogenic after transplantation. *Stem Cells* 24, 1054–1062 (2004).

- **Article**

274. Jendelova, P. *et al.* Magnetic resonance tracking of transplanted bone marrow and embryonic stem cell-derived nanoparticles in rat brain and spinal cord. *J. Neurosci. Res.* 76, 232–243 (2004).

- **ChemPort**

275. Shier, D. N. *Hole's Human Anatomy and Physiology*. 7th edn (TM Higher Education Group, 2004).

Author affiliations

1. Centre for the Cellular Basis of Behaviour, MRC Centre for Neurodegeneration Research, King's College London, P.O. Box 39, 1–2 WW Ground, Denmark Hill, London SE5 8AF, UK.
2. Wolfson Centre for Age-Related Diseases, King's College London, 16–18 Newcomen Street, London SE1 1UL, UK.
3. Laboratory of Genetics, The Salk Institute for Biological Studies, La Jolla, California 92037, USA.
4. These authors contributed equally to this work.

Correspondence to: Fred H. Gage³ Email: gage@salk.edu

MORE ARTICLES LIKE THIS

These links to content published by NPG are automatically generated.

NEWS AND VIEWS

Just one word: plasticity

Nature Neuroscience News and Views (01 Mar 2004)

Stem cells in the injured spinal cord: reducing the pain and increasing the gain

Nature Neuroscience News and Views (01 Mar 2005)

RESEARCH

cAMP and Schwann cells promote axonal growth and functional recovery after spinal cord injury

Nature Medicine Article (01 Jun 2004)

See all 9 matches for Research

- [**Previous article**](#)
- [**Next article**](#)

Top

Nature Reviews Neuroscience

ISSN: 1471-003X

EISSN: 1471-0048

- [**About NPG**](#)
 - [**Contact NPG**](#)
 - [**Nature jobs.com**](#)
 - [**Privacy policy**](#)
 - [**Legal notice**](#)
 - [**Accessibility statement**](#)
 - [**RSS web feeds**](#)
 - [**Help**](#)
-

© 2007 Nature Publishing Group – partner of AGORA, HINARI, CrossRef and COUNTER